

FORMULA BOOK STATISTICS CA FOUNDATION DEC 2023

CA. PRANAV POPAT

FORMULA MARATHON MATHS
SESSION LINK:

<https://www.youtube.com/live/OqUu7X8TYMQ?si=IPq9Bwh-xJYbRGwb>

JOIN TELEGRAM CHANNEL FOR ALL
UPDATES AND NOTES:

<https://telegram.me/learnwithpranav>



Formula 1	Class Boundary				
	Mutually Exclusive Classification		UCB = UCL and LCB = LCL		
	Mutually Inclusive Classification		UCB = UCL + 0.5 and LCB = LCL - 0.5		
Formula 2	Mid-Point / Class Mark of Class Interval: $\frac{LCL + UCL}{2}$ or $\frac{LCB + UCB}{2}$				
Formula 3	Class Length / Width of Class / Size of Class: UCB – LCB				
Formula 4	Frequency Density of a Class: $\frac{\text{Frequency of the class}}{\text{Class length of the class}}$				
Formula 5	Relative Frequency: $\frac{\text{Frequency of the class}}{\text{Total Frequency of distribution}}$				
	Percentage Frequency: $\frac{\text{Frequency of the class}}{\text{Total Frequency of distribution}} \times 100$				
Formula 6	AM of Discrete Distribution/Series: $\bar{x} = \frac{x_1 + x_2 + x_3 + \dots + x_n}{n}$ in short $\bar{x} = \frac{\sum x}{n}$				
Formula 7	AM of Frequency Distribution: $\bar{x} = \frac{\sum fx}{N}$				
	In case of ungrouped distribution		x = individual value		
	In case of grouped frequency distribution		x = mid-point of class interval		
Formula 8	AM using assumed mean / step deviation method $\bar{x} = A + \frac{\sum fd}{N} \times C$ where $d = \frac{x - A}{C}$, A is assumed mean, C is class length				
Formula 9	The algebraic sum of deviations of a set of observations from their AM is zero $\sum(x - \bar{x}) = 0$				
Formula 10	Combined AM: $\bar{x}_c = \frac{n_1\bar{x}_1 + n_2\bar{x}_2}{n_1 + n_2}$				
Formula 11	Median in case of discrete distribution				
	If number of observations are odd		Median is middle term		
	If number of observations are even		AM of two middle terms		
Same formula is used for ungrouped frequency distribution					
Formula 12	Median in case of grouped frequency distribution				
	Step 1	Prepare a less than type cumulative frequency distribution			
	Step 2	Calculate $\frac{N}{2}$ and check between which class boundaries it falls and call it as Median Class			
	Step 3	I_1	N_u	N_l	C
		LCB of Median Class	Cum Freq. of Median Class	Cum. Freq. of Pre-Median Class	Class length of Median Class
Step 4	Apply Formula $Me = I_1 + \left(\frac{\frac{N}{2} - N_l}{N_u - N_l} \right) \times C$				



Formula 13	For a set of observations, the sum of absolute deviations is minimum, when the deviations are taken from the median. $\sum(x - \bar{x}) = 0$ is minimum							
Formula 14	Quartiles in case of discrete observations:							
	<table border="1"> <thead> <tr> <th>First Quartile</th> <th>Second Quartile</th> <th>Third Quartile</th> </tr> </thead> <tbody> <tr> <td>$Q_1 = \left((n+1) \times \frac{1}{4} \right)^{\text{th}}$ term</td> <td>$Q_2 = \left((n+1) \times \frac{2}{4} \right)^{\text{th}}$ term</td> <td>$Q_3 = \left((n+1) \times \frac{3}{4} \right)^{\text{th}}$ term</td> </tr> </tbody> </table>	First Quartile	Second Quartile	Third Quartile	$Q_1 = \left((n+1) \times \frac{1}{4} \right)^{\text{th}}$ term	$Q_2 = \left((n+1) \times \frac{2}{4} \right)^{\text{th}}$ term	$Q_3 = \left((n+1) \times \frac{3}{4} \right)^{\text{th}}$ term	
First Quartile	Second Quartile	Third Quartile						
$Q_1 = \left((n+1) \times \frac{1}{4} \right)^{\text{th}}$ term	$Q_2 = \left((n+1) \times \frac{2}{4} \right)^{\text{th}}$ term	$Q_3 = \left((n+1) \times \frac{3}{4} \right)^{\text{th}}$ term						
Note: above formula gives the term. Final value to be calculated based on the term								
Formula 15	Deciles in case of discrete observations:							
	<table border="1"> <thead> <tr> <th>First Decile</th> <th>Second Decile</th> <th>Ninth Decile</th> </tr> </thead> <tbody> <tr> <td>$D_1 = \left((n+1) \times \frac{1}{10} \right)^{\text{th}}$ term</td> <td>$D_2 = \left((n+1) \times \frac{2}{10} \right)^{\text{th}}$ term</td> <td>$D_9 = \left((n+1) \times \frac{9}{10} \right)^{\text{th}}$ term</td> </tr> </tbody> </table>	First Decile	Second Decile	Ninth Decile	$D_1 = \left((n+1) \times \frac{1}{10} \right)^{\text{th}}$ term	$D_2 = \left((n+1) \times \frac{2}{10} \right)^{\text{th}}$ term	$D_9 = \left((n+1) \times \frac{9}{10} \right)^{\text{th}}$ term	
First Decile	Second Decile	Ninth Decile						
$D_1 = \left((n+1) \times \frac{1}{10} \right)^{\text{th}}$ term	$D_2 = \left((n+1) \times \frac{2}{10} \right)^{\text{th}}$ term	$D_9 = \left((n+1) \times \frac{9}{10} \right)^{\text{th}}$ term						
Note: above formula gives the term. Final value to be calculated based on the term								
Formula 16	Percentiles in case of discrete observations:							
	<table border="1"> <thead> <tr> <th>First Percentile</th> <th>Second Percentile</th> <th>99th Percentile</th> </tr> </thead> <tbody> <tr> <td>$P_1 = \left((n+1) \times \frac{1}{100} \right)^{\text{th}}$ term</td> <td>$P_2 = \left((n+1) \times \frac{2}{100} \right)^{\text{th}}$ term</td> <td>$P_{99} = \left((n+1) \times \frac{99}{100} \right)^{\text{th}}$ term</td> </tr> </tbody> </table>	First Percentile	Second Percentile	99 th Percentile	$P_1 = \left((n+1) \times \frac{1}{100} \right)^{\text{th}}$ term	$P_2 = \left((n+1) \times \frac{2}{100} \right)^{\text{th}}$ term	$P_{99} = \left((n+1) \times \frac{99}{100} \right)^{\text{th}}$ term	
First Percentile	Second Percentile	99 th Percentile						
$P_1 = \left((n+1) \times \frac{1}{100} \right)^{\text{th}}$ term	$P_2 = \left((n+1) \times \frac{2}{100} \right)^{\text{th}}$ term	$P_{99} = \left((n+1) \times \frac{99}{100} \right)^{\text{th}}$ term						
Note: above formula gives the term. Final value to be calculated based on the term								
Formula 17	Quartiles in case of Grouped Frequency Distribution: Steps are like median with few modifications.							
	<table border="1"> <thead> <tr> <th>1st Quartile</th> <th>3rd Quartile</th> </tr> </thead> <tbody> <tr> <td>Find Q_1 class using $\frac{N}{4}$</td> <td>Find Q_3 class using $\frac{3N}{4}$</td> </tr> <tr> <td>$Q_1 = l_1 + \left(\frac{\frac{N}{4} - N_l}{N_u - N_l} \right) \times C$</td> <td>$Q_3 = l_1 + \left(\frac{\frac{3N}{4} - N_l}{N_u - N_l} \right) \times C$</td> </tr> </tbody> </table>	1 st Quartile	3 rd Quartile	Find Q_1 class using $\frac{N}{4}$	Find Q_3 class using $\frac{3N}{4}$	$Q_1 = l_1 + \left(\frac{\frac{N}{4} - N_l}{N_u - N_l} \right) \times C$	$Q_3 = l_1 + \left(\frac{\frac{3N}{4} - N_l}{N_u - N_l} \right) \times C$	
1 st Quartile	3 rd Quartile							
Find Q_1 class using $\frac{N}{4}$	Find Q_3 class using $\frac{3N}{4}$							
$Q_1 = l_1 + \left(\frac{\frac{N}{4} - N_l}{N_u - N_l} \right) \times C$	$Q_3 = l_1 + \left(\frac{\frac{3N}{4} - N_l}{N_u - N_l} \right) \times C$							
Formula 18	Deciles in case of Grouped Frequency Distribution: Steps are like median with few modifications.							
	<table border="1"> <thead> <tr> <th>1st Decile</th> <th>9th Decile</th> </tr> </thead> <tbody> <tr> <td>Find D_1 class using $\frac{N}{10}$</td> <td>Find D_9 class using $\frac{9N}{10}$</td> </tr> <tr> <td>$D_1 = l_1 + \left(\frac{\frac{N}{10} - N_l}{N_u - N_l} \right) \times C$</td> <td>$D_9 = l_1 + \left(\frac{\frac{9N}{10} - N_l}{N_u - N_l} \right) \times C$</td> </tr> </tbody> </table>	1 st Decile	9 th Decile	Find D_1 class using $\frac{N}{10}$	Find D_9 class using $\frac{9N}{10}$	$D_1 = l_1 + \left(\frac{\frac{N}{10} - N_l}{N_u - N_l} \right) \times C$	$D_9 = l_1 + \left(\frac{\frac{9N}{10} - N_l}{N_u - N_l} \right) \times C$	
1 st Decile	9 th Decile							
Find D_1 class using $\frac{N}{10}$	Find D_9 class using $\frac{9N}{10}$							
$D_1 = l_1 + \left(\frac{\frac{N}{10} - N_l}{N_u - N_l} \right) \times C$	$D_9 = l_1 + \left(\frac{\frac{9N}{10} - N_l}{N_u - N_l} \right) \times C$							
Formula 19	Percentiles in case of Grouped Frequency Distribution: Steps are like median with few modifications.							
	<table border="1"> <thead> <tr> <th>1st Percentile</th> <th>99th Percentile</th> </tr> </thead> <tbody> <tr> <td>Find P_1 class using $\frac{N}{100}$</td> <td>Find P_{99} class using $\frac{99N}{100}$</td> </tr> </tbody> </table>	1 st Percentile	99 th Percentile	Find P_1 class using $\frac{N}{100}$	Find P_{99} class using $\frac{99N}{100}$			
1 st Percentile	99 th Percentile							
Find P_1 class using $\frac{N}{100}$	Find P_{99} class using $\frac{99N}{100}$							



	$P_1 = l_1 + \left(\frac{\frac{N}{100} - N_i}{N_u - N_i} \right) \times C$ $P_{99} = l_1 + \left(\frac{\frac{99N}{10} - N_i}{N_u - N_i} \right) \times C$								
Formula 20	<p>Mode in case of discrete observation: observation repeating for maximum no. of times or observation with highest frequency Note: There can be multiple modes also. If all observations are having same frequency, then there is no mode.</p>								
Formula 21	<p>Mode in case of grouped frequency distribution: Find Modal Class (Class with highest frequency) then apply below formula</p> $Mo = l_1 + \left(\frac{f_0 - f_{-1}}{2f_0 - f_{-1} - f_1} \right) \times C$ <p>where, l_1 = LCB of modal class f_0 = frequency of modal class, f_{-1} = frequency of pre-modal class, f_1 = frequency of post modal class, C = class length of modal class</p>								
Formula 22	<p>Relationship between Mean, Median and Mode in case of Symmetrical Distribution: Mean = Median = Mode</p>								
Formula 23	<p>Relationship between Mean, Median and Mode in case of moderately skewed distribution: Mean – Mode = 3 (Mean – Median)</p>								
Formula 24	<p>Geometric Mean in case of discrete positive observations:</p> $G = (x_1 \times x_2 \times \dots \times x_n)^{1/n}$								
Formula 25	<p>Geometric Mean in case of frequency distribution:</p> $G = (x_1^{f_1} \times x_2^{f_2} \times \dots \times x_n^{f_n})^{1/N}$								
Formula 26	<p>Harmonic Mean in case of discrete observations: $H = \frac{n}{\sum(\frac{1}{x})}$</p>								
Formula 27	<p>Harmonic Mean in case of frequency distribution: $H = \frac{N}{\sum(\frac{f}{x})}$</p>								
Formula 28	<p>Combined HM = $\frac{\frac{n_1 + n_2}{\frac{n_1}{H_1} + \frac{n_2}{H_2}}}$</p>								
Formula 29	<p>Relationship between AM, GM and HM</p> <table border="1"> <thead> <tr> <th>Situation</th> <th>Relationship</th> </tr> </thead> <tbody> <tr> <td>When all the observations are identical / same</td> <td>AM = GM = HM</td> </tr> <tr> <td>When all the observations are distinct / different</td> <td>AM > GM > HM</td> </tr> <tr> <td>In General</td> <td>AM ≥ GM ≥ HM</td> </tr> </tbody> </table>	Situation	Relationship	When all the observations are identical / same	AM = GM = HM	When all the observations are distinct / different	AM > GM > HM	In General	AM ≥ GM ≥ HM
Situation	Relationship								
When all the observations are identical / same	AM = GM = HM								
When all the observations are distinct / different	AM > GM > HM								
In General	AM ≥ GM ≥ HM								
Formula 30	<p>Range in case of discrete observations: L – S where L = Largest Observation, S = Smallest Observation</p>								
Formula 31	<p>Range in case of Grouped Frequency Distribution: L – S L = UCB of last class interval, S = LCB of first-class interval</p>								
Formula 32	<p>Coefficient of Range $\frac{L - S}{L + S} \times 100$</p>								
Formula 33	<p>Mean Deviation in case of discrete observations</p>								



	$MD_A = \frac{1}{n} \sum x - A $ where A is any appropriate central tendency (as given)
Formula 34	Mean Deviation (in case of grouped frequency distributions) $MD_A = \frac{1}{N} \sum f x - A $ where A is any appropriate central tendency (as given)
Formula 35	Coefficient of Mean Deviation: $\frac{\text{Mean Deviation about A}}{A} \times 100$
Formula 36	Standard Deviation in case of discrete observations: $\sigma_x = SD_x = \sqrt{\frac{\sum (x - \bar{x})^2}{n}}$ or shorter formula $\sigma_x = SD_x = \sqrt{\frac{\sum x^2}{n} - (\bar{x})^2}$
Formula 37	Standard Deviation in case of grouped frequency observations $\sigma_x = SD_x = \sqrt{\frac{\sum f(x - \bar{x})^2}{N}}$ or shorter formula $\sigma_x = SD_x = \sqrt{\frac{\sum fx^2}{N} - (\bar{x})^2}$
Formula 38	Coefficient of Variation: $\frac{SD_x}{\bar{x}} \times 100$
Formula 39	If there are only two observations, then SD is half of range $SD = \frac{ a - b }{2}$
Formula 40	Standard Deviation of first n natural numbers: $s = \sqrt{\frac{n^2 - 1}{12}}$
Formula 41	Combined SD: $SD_c = \sqrt{\frac{n_1 s_1^2 + n_2 s_2^2 + n_1 d_1^2 + n_2 d_2^2}{n_1 + n_2}}$ $d_1 = \bar{x}_c - \bar{x}_1$ and $d_2 = \bar{x}_c - \bar{x}_2$
Formula 42	If all the observations are constant, then SD/ MD/ Range is ZERO
Formula 43	Change of Origin and Scale: No effect of change of origin but affected by change of scale in the magnitude (ignore sign) $SD_y = b SD_x$ Note: same thing will apply to all the measures of dispersion
Formula 44	Quartile Deviation: $QD_x = \frac{Q_3 - Q_1}{2}$
Formula 45	Coefficient of Quartile Deviation: $\frac{Q_3 - Q_1}{Q_3 + Q_1} \times 100$
Formula 46	Relationship between SD, MD and QD $4SD = 5MD = 6QD$ or $SD : MD : QD = 15 : 12 : 10$
Formula 47	Basic Formula of Probability: $P(A) = \frac{\text{No. of favorable events to A}}{\text{Total no. of events}}$
Formula 48	Odds in favour of Event A: $\frac{\text{no. of favorable events}}{\text{no. of unfavorable events}}$
Formula 49	Odds against an Event A: $\frac{\text{no. of unfavorable events}}{\text{no. of favorable events}}$
Formula 50	Number of total outcomes of a random experiment: If an experiment results in p outcomes and if it is repeated q times, then Total number of outcomes is p^q



Formula 51	Relative Frequency Probability $\frac{\text{no. of times the event occurred during experimental trials}}{\text{total no. of trials}} = \frac{f_A}{n}$						
Formula 52	Set Based Probability: $P(A) = \frac{\text{no. of sample points in A}}{\text{no. of sample points in S}} = \frac{n(A)}{n(S)}$ here A is Event Set and S is Sample Space						
Formula 53	Addition Theorem 1: In case of two mutually exclusive events A and B $P(A \cup B) = P(A+B) = P(A \text{ or } B) = P(A) + P(B)$						
Formula 54	Addition Theorem 2: In case of two or more mutually exclusive events $P(A_1 \cup A_2 \cup A_3 \cup \dots) = P(A_1) + P(A_2) + P(A_3) + \dots$						
Formula 55	Addition Theorem 3: For any two events $P(A \cup B) = P(A) + P(B) - P(A \cap B)$						
Formula 56	Addition Theorem 4: In case of any three events $P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(B \cap C) - P(A \cap C) + P(A \cap B \cap C)$						
Formula 57	Conditional Probability of Event B when Event A is already occurred $P(B/A) = \frac{P(B \cap A)}{P(A)}$ provided $P(A) \neq 0$						
Formula 58	Conditional Probability of Event A when Event B is already occurred $P(A/B) = \frac{P(B \cap A)}{P(B)}$ provided $P(B) \neq 0$						
Formula 59	Compound Theorem: In case of two dependent events $P(A \cap B) = P(B) \times P(A/B)$ or $P(A \cap B) = P(A) \times P(B/A)$						
Formula 60	Compound Theorem: In case of two independent events $P(A \cap B) = P(A) \times P(B)$						
Formula 61	Expected value of a Probability Distribution: $E(x) = \sum p_i x_i$ Also, $E(x) = \mu$ (here μ means mean of probability distribution)						
Formula 62	Variance of Probability Distribution: $V(x) = E(x - \mu)^2 = E(x^2) - [E(x)]^2$						
Formula 63	Probability Mass Function in case of Binomial Distribution: $f(x) = P(X=x) = {}^n C_x p^x q^{n-x}$						
Formula 64	Mean of Binomial Distribution: $\mu = np$ Variance of Binomial Distribution: $\sigma^2 = npq$						
Formula 65	Mode in case of Binomial Distribution: <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 20%;">Step 1</td> <td>Calculate $(n+1)p$</td> </tr> <tr> <td>Step 2A</td> <td>If $(n+1)p$ is an integer, there will be two modes: $\mu_0 = (n+1)p$ & $[(n+1)p - 1]$</td> </tr> <tr> <td>Step 2B</td> <td>If $(n+1)p$ is a non-integer, there will be only one mode: $\mu_0 =$ largest integer contained in $(n+1)p$</td> </tr> </table>	Step 1	Calculate $(n+1)p$	Step 2A	If $(n+1)p$ is an integer, there will be two modes: $\mu_0 = (n+1)p$ & $[(n+1)p - 1]$	Step 2B	If $(n+1)p$ is a non-integer, there will be only one mode: $\mu_0 =$ largest integer contained in $(n+1)p$
Step 1	Calculate $(n+1)p$						
Step 2A	If $(n+1)p$ is an integer, there will be two modes: $\mu_0 = (n+1)p$ & $[(n+1)p - 1]$						
Step 2B	If $(n+1)p$ is a non-integer, there will be only one mode: $\mu_0 =$ largest integer contained in $(n+1)p$						
Formula 66	Probability Mass Function in case of Poisson Distribution: $f(x) = P(X=x) = \frac{e^{-m} m^x}{x!}$						
Formula 67	Mean of Poisson Distribution: $\mu = m$ Variance of Poisson Distribution: $\sigma^2 = m$						



	SD of Poisson Distribution: $\sigma = \sqrt{m}$															
Formula 68	Mode in case of Poisson Distribution:															
	If m is an integer there will be two modes: $\mu_0 = m$ & $m-1$															
	If m is a non-integer there will be only one mode: largest integer contained in m															
Formula 69	Probability Density Function in case of Normal Distribution $f(x) = \frac{1}{\sigma\sqrt{2\pi}} \cdot e^{-\left(\frac{x-\mu}{\sigma}\right)^2 \frac{1}{2}}$															
Formula 70	Mean Deviation in case of Normal Distribution: MD = 0.8σ															
Formula 71	Quartiles in case of Normal Distribution: $Q_1 = \mu - 0.675\sigma$ & $Q_3 = \mu + 0.675\sigma$															
Formula 72	Quartile Deviation in case of Normal Distribution: QD = 0.675σ															
Formula 73	Points of Inflex of Normal Curve: $\mu - \sigma$ & $\mu + \sigma$															
Formula 74	In case of Normal Distribution, Ratio between QD: MD: SD = 10:12:15															
Formula 75	Conditions of Standard Normal Distribution: Mean = 0, SD = 1															
Formula 76	Z Score: $Z = \frac{(x - \mu)}{\sigma}$															
Formula 77	Area under Normal Curve (Popular Intervals)															
	<table border="1"> <thead> <tr> <th>From</th> <th>To</th> <th>Area under Normal Curve Probability</th> </tr> </thead> <tbody> <tr> <td>μ</td> <td>$\mu + \sigma$</td> <td>34.135%</td> </tr> <tr> <td>$\mu + \sigma$</td> <td>$\mu + 2\sigma$</td> <td>13.59%</td> </tr> <tr> <td>$\mu + 2\sigma$</td> <td>$\mu + 3\sigma$</td> <td>2.14%</td> </tr> <tr> <td>$\mu + 3\sigma$</td> <td>$+\infty$</td> <td>0.135%</td> </tr> </tbody> </table>	From	To	Area under Normal Curve Probability	μ	$\mu + \sigma$	34.135%	$\mu + \sigma$	$\mu + 2\sigma$	13.59%	$\mu + 2\sigma$	$\mu + 3\sigma$	2.14%	$\mu + 3\sigma$	$+\infty$	0.135%
	From	To	Area under Normal Curve Probability													
	μ	$\mu + \sigma$	34.135%													
	$\mu + \sigma$	$\mu + 2\sigma$	13.59%													
$\mu + 2\sigma$	$\mu + 3\sigma$	2.14%														
$\mu + 3\sigma$	$+\infty$	0.135%														
Formula 78	For a $p \times q$ bivariate frequency distribution:															
	<table border="1"> <tr> <td>Number of cells</td> <td>pq</td> </tr> <tr> <td>Number of marginal distributions</td> <td>2</td> </tr> <tr> <td>Number of conditional distributions</td> <td>$p+q$</td> </tr> </table>	Number of cells	pq	Number of marginal distributions	2	Number of conditional distributions	$p+q$									
	Number of cells	pq														
Number of marginal distributions	2															
Number of conditional distributions	$p+q$															
Formula 79	Karl Pearson's Product Moment Correlation Coefficient: $r_{xy} = \frac{\text{Cov}(x, y)}{(\sigma_x \times \sigma_y)}$															
Formula 80	Covariance between two variables: $\text{Cov}(x, y) = \frac{\sum(x - \bar{x})(y - \bar{y})}{n}$ or $\frac{\sum xy}{n} - \bar{x} \cdot \bar{y}$															
Formula 81	Spearman's Rank Correlation Coefficient: $r_R = 1 - \frac{6\sum d^2}{n(n^2 - 1)}$ here d means difference in ranks of both variables															
Formula 82	Spearman's Rank Correlation Coefficient (in case of tied values)															
	$r_R = 1 - \frac{6(\sum d^2 + A)}{n(n^2 - 1)}$ here A is adjustment value															
	$A = \frac{\sum(t^3 - t)}{12}$ where t = tie length (calculate t value for each of the ties)															
Formula 83	Coefficient of Concurrent Deviations															



	$r_c = \pm \sqrt{\pm \left(\frac{2c-m}{m} \right)}$ <p>where c is number of concurrent deviations (same direction) m is number of pairs compared (equals to n-1)</p>
Formula 84	<p>Regression Coefficients:</p> <p>Y on X: $b_{yx} = r \cdot \frac{SD_y}{SD_x}$ or $b_{yx} = \frac{cov(x,y)}{(SD_x)^2}$</p> <p>X on Y: $b_{xy} = r \cdot \frac{SD_x}{SD_y}$ or $b_{xy} = \frac{cov(x,y)}{(SD_y)^2}$</p>
Formula 85	<p>Correlation Coefficient is the GM of regression coefficients:</p> $r_{xy} = \pm \sqrt{b_{xy} \times b_{yx}}$ <p>Note: r_{xy}, b_{xy}, b_{yx} all will have same sign</p>
Formula 86	<p>Change of Origin/ Scale for Regression Coefficients: Origin no impact, Scale impact of both magnitude and sign.</p> $b_{vu} = b_{yx} \times \frac{\text{change of scale of } y}{\text{change of scale of } x}$ $b_{uv} = b_{xy} \times \frac{\text{change of scale of } x}{\text{change of scale of } y}$
Formula 87	Two regression lines (if not identical) will intersect at the point (\bar{x}, \bar{y})
Formula 88	<p>Coefficient of Determination/ Explained Variance/ Accounted Variance:</p> $(r_{xy})^2$
Formula 89	<p>Coefficient of Non-determination/ Un-explained Variance/ Un-accounted Variance:</p> $1 - (r_{xy})^2$
Formula 90	Probable Error in correlation: $0.6745 \times \frac{1-r^2}{\sqrt{N}}$
Formula 91	Error Limits of Population Correlation Coefficient: $r \pm PE$
Formula 92	Price Relatives: $\frac{P_n}{P_0}$, Quantity Relatives: $\frac{Q_n}{Q_0}$, Value Relatives: $\frac{V_n}{V_0}$
Formula 93	Simple Aggregative Index: $\frac{\sum P_n}{\sum P_0} \times 100$
Formula 94	Simple Average of Relatives – Method Index: $\frac{\sum \frac{P_n}{P_0}}{n}$
Formula 95	<p>Laspeyres Index (weight – base year quantity weight)</p> $\frac{\sum P_n Q_0}{\sum P_0 Q_0} \times 100$
Formula 96	<p>Paasche's Index (weight – current year quantity weight)</p> $\frac{\sum P_n Q_n}{\sum P_0 Q_n} \times 100$
Formula 97	Marshall-Edgeworth Index (weight – sum of both current and base quantity)



	$\frac{\sum P_n (Q_0 + Q_n)}{\sum P_0 (Q_0 + Q_n)} \times 100$
Formula 98	Fisher's Ideal Index: GM of Laspeyres Index and Paasche's Index $\sqrt{\frac{\sum P_n Q_0}{\sum P_0 Q_0} \times \frac{\sum P_n Q_n}{\sum P_0 Q_n}} \times 100$
Formula 99	Bowley's Index: AM of Laspeyres Index and Paasche's Index $\frac{\frac{\sum P_n Q_0}{\sum P_0 Q_0} + \frac{\sum P_n Q_n}{\sum P_0 Q_n}}{2} \times 100$

About CA. Pranav Popat Sir

- He is a Chartered Accountant (Inter and Final Both Groups in First Attempt) with 6+ years of experience.
- He is an Educator by Passion and his Choice (Dil Se ❤️)
- Taught 10k + students
- He teaches subjects of Maths, LR and Stats (Paper 3) at CA Foundation Level and Cost & Management Accounting (Paper 3) at CA Intermediate Level.

Hope this formula book helps you in revising all formulas and become helpful to you during exam time, I made this with my whole heart, make best use of it and I just want one thing in return – share these notes to every student who really needs this.

Wishing you ALL THE BEST for upcoming examinations, see you soon in Inter Costing!!!

Ab mushkil nahi kuch bhi, nahi kuch bhi!!!

With Lots of Love

CA. Pranav Popat (P^2 SIR)

