

Stats

24 May 2024 09:50



Stats
Formula...

FORMULA BOOK STATISTICS CA FOUNDATION JUNE 2024

CA. PRANAV POPAT

FORMULA MARATHON STATS
SESSION LINK:

<https://www.youtube.com/watch?v=XXv5wRqso7w>

JOIN TELEGRAM CHANNEL FOR ALL
UPDATES AND NOTES:

<https://telegram.me/learnwithpranav>



Formula 1	Class Boundary	
	Mutually Exclusive Classification 10-20 20-30 30-40	UCB = UCL LCB = LCL
	Mutually Inclusive Classification 10-19 20-29 30-39	UCB = UCL + 0.5 LCB = LCL - 0.5
Formula 2	Mid-Point / Class Mark of Class Interval: $\frac{LCL + UCL}{2}$ or $\frac{LCB + UCB}{2}$	
Formula 3	Class Length / Width of Class / Size of Class: UCB - LCB only	
Formula 4	Frequency Density of a Class: $\frac{\text{class frequency}}{\text{class length}}$	
Formula 5	Relative Frequency: $\frac{\text{class frequency}}{\text{Total frequency}}$	
	Percentage Frequency: $\frac{\text{class freq}}{\text{Total freq}} \times 100$	
Formula 6	AM of Discrete Distribution/Series: $\frac{x_1 + x_2 + x_3 + \dots + x_n}{n} = \frac{\sum x}{n}$	
Formula 7	AM of Frequency Distribution: $\frac{\sum fx}{N}$ $N = \sum f$	
	In case of ungrouped distribution	x = individual value
	In case of grouped frequency distribution	x = mid-point of class interval
Formula 8	AM using assumed mean / step deviation method $A + \frac{\sum fd}{N} \times C$ where $d = \frac{x-A}{C}$, A is assumed mean, C is class length	
Formula 9	The algebraic sum of deviations of a set of observations from their AM is <u>zero</u>	



Formula 10	Combined AM: $\bar{x}_c = \frac{n_1\bar{x}_1 + n_2\bar{x}_2}{n_1 + n_2}$									
Formula 11	Median in case of discrete distribution <i>arrange data in ascending order</i>									
	If number of observations are odd	Median is <u>middle term</u>								
	If number of observations are even	Median is <u>avg of two middle terms</u>								
Same formula is used for ungrouped frequency distribution										
Formula 12	Median in case of grouped frequency distribution									
	Step 1	Prepare a less than type cumulative frequency distribution								
	Step 2	Calculate $\frac{N}{2}$ and check between which class boundaries it falls and call it as Median Class								
	Step 3	<table border="1"> <thead> <tr> <th>l_1</th> <th>N_u</th> <th>N_l</th> <th>C</th> </tr> </thead> <tbody> <tr> <td>LCB of Median Class</td> <td>Cum Freq. of Median Class</td> <td>Cum. Freq. of Pre-Median Class</td> <td>Class length of Median Class</td> </tr> </tbody> </table>	l_1	N_u	N_l	C	LCB of Median Class	Cum Freq. of Median Class	Cum. Freq. of Pre-Median Class	Class length of Median Class
l_1	N_u	N_l	C							
LCB of Median Class	Cum Freq. of Median Class	Cum. Freq. of Pre-Median Class	Class length of Median Class							
Step 4	Apply Formula $\text{Median} = l_1 + \left(\frac{\frac{N}{2} - N_l}{N_u - N_l} \right) \times C$		$N_u - N_l = f$							
Formula 13	For a set of observations, the sum of absolute deviations is <u>minimum</u> when the deviations are taken from the median. $\sum(x - \bar{x}) = 0$ is $\sum x - \text{Median} $ is minimum									
Formula 14	Quartiles in case of discrete observations:									
	First Quartile $Q_1 = \left((n+1) \times \frac{1}{4} \right)^{\text{th}}$ term	Second Quartile $Q_2 = \left((n+1) \times \frac{2}{4} \right)^{\text{th}}$ term	Third Quartile $Q_3 = \left((n+1) \times \frac{3}{4} \right)^{\text{th}}$ term							

Rank = $(n+1) \times p^{\text{th}}$ term



	Note: above formula gives the <u>term</u> . <u>Final value</u> to be calculated based on the term							
Formula 15	Deciles in case of discrete observations:							
	<table border="1"> <thead> <tr> <th>First Decile</th> <th>Second Decile</th> <th>Ninth Decile</th> </tr> </thead> <tbody> <tr> <td>$D_1 = \left((n+1) \times \frac{1}{10} \right)^{\text{th}}$ term</td> <td>$D_2 = \left((n+1) \times \frac{2}{10} \right)^{\text{th}}$ term</td> <td>$D_9 = \left((n+1) \times \frac{9}{10} \right)^{\text{th}}$ term</td> </tr> </tbody> </table>	First Decile	Second Decile	Ninth Decile	$D_1 = \left((n+1) \times \frac{1}{10} \right)^{\text{th}}$ term	$D_2 = \left((n+1) \times \frac{2}{10} \right)^{\text{th}}$ term	$D_9 = \left((n+1) \times \frac{9}{10} \right)^{\text{th}}$ term	
First Decile	Second Decile	Ninth Decile						
$D_1 = \left((n+1) \times \frac{1}{10} \right)^{\text{th}}$ term	$D_2 = \left((n+1) \times \frac{2}{10} \right)^{\text{th}}$ term	$D_9 = \left((n+1) \times \frac{9}{10} \right)^{\text{th}}$ term						
	Note: above formula gives the term. Final value to be calculated based on the term							
Formula 16	Percentiles in case of discrete observations:							
	<table border="1"> <thead> <tr> <th>First Percentile</th> <th>Second Percentile</th> <th>99th Percentile</th> </tr> </thead> <tbody> <tr> <td>$P_1 = \left((n+1) \times \frac{1}{100} \right)^{\text{th}}$ term</td> <td>$P_2 = \left((n+1) \times \frac{2}{100} \right)^{\text{th}}$ term</td> <td>$P_{99} = \left((n+1) \times \frac{99}{100} \right)^{\text{th}}$ term</td> </tr> </tbody> </table>	First Percentile	Second Percentile	99 th Percentile	$P_1 = \left((n+1) \times \frac{1}{100} \right)^{\text{th}}$ term	$P_2 = \left((n+1) \times \frac{2}{100} \right)^{\text{th}}$ term	$P_{99} = \left((n+1) \times \frac{99}{100} \right)^{\text{th}}$ term	
First Percentile	Second Percentile	99 th Percentile						
$P_1 = \left((n+1) \times \frac{1}{100} \right)^{\text{th}}$ term	$P_2 = \left((n+1) \times \frac{2}{100} \right)^{\text{th}}$ term	$P_{99} = \left((n+1) \times \frac{99}{100} \right)^{\text{th}}$ term						
	Note: above formula gives the term. Final value to be calculated based on the term							
Formula 17	Quartiles in case of Grouped Frequency Distribution: Steps are like median with few modifications.							
	<table border="1"> <thead> <tr> <th>1st Quartile</th> <th>3rd Quartile</th> </tr> </thead> <tbody> <tr> <td>Find Q_1 class using $\frac{N}{4}$</td> <td>Find Q_3 class using $\frac{3N}{4}$</td> </tr> <tr> <td>$Q_1 = l_1 + \left(\frac{\frac{N}{4} - N_l}{N_u - N_l} \right) \times C$</td> <td>$Q_3 = l_1 + \left(\frac{\frac{3N}{4} - N_l}{N_u - N_l} \right) \times C$</td> </tr> </tbody> </table>	1 st Quartile	3 rd Quartile	Find Q_1 class using $\frac{N}{4}$	Find Q_3 class using $\frac{3N}{4}$	$Q_1 = l_1 + \left(\frac{\frac{N}{4} - N_l}{N_u - N_l} \right) \times C$	$Q_3 = l_1 + \left(\frac{\frac{3N}{4} - N_l}{N_u - N_l} \right) \times C$	
1 st Quartile	3 rd Quartile							
Find Q_1 class using $\frac{N}{4}$	Find Q_3 class using $\frac{3N}{4}$							
$Q_1 = l_1 + \left(\frac{\frac{N}{4} - N_l}{N_u - N_l} \right) \times C$	$Q_3 = l_1 + \left(\frac{\frac{3N}{4} - N_l}{N_u - N_l} \right) \times C$							
Formula 18	Deciles in case of Grouped Frequency Distribution: Steps are like median with few modifications.							
	<table border="1"> <thead> <tr> <th>1st Decile</th> <th>9th Decile</th> </tr> </thead> <tbody> <tr> <td>Find D_1 class using $\frac{N}{10}$</td> <td>Find D_9 class using $\frac{9N}{10}$</td> </tr> <tr> <td>$D_1 = l_1 + \left(\frac{\frac{N}{10} - N_l}{N_u - N_l} \right) \times C$</td> <td>$D_9 = l_1 + \left(\frac{\frac{9N}{10} - N_l}{N_u - N_l} \right) \times C$</td> </tr> </tbody> </table>	1 st Decile	9 th Decile	Find D_1 class using $\frac{N}{10}$	Find D_9 class using $\frac{9N}{10}$	$D_1 = l_1 + \left(\frac{\frac{N}{10} - N_l}{N_u - N_l} \right) \times C$	$D_9 = l_1 + \left(\frac{\frac{9N}{10} - N_l}{N_u - N_l} \right) \times C$	
1 st Decile	9 th Decile							
Find D_1 class using $\frac{N}{10}$	Find D_9 class using $\frac{9N}{10}$							
$D_1 = l_1 + \left(\frac{\frac{N}{10} - N_l}{N_u - N_l} \right) \times C$	$D_9 = l_1 + \left(\frac{\frac{9N}{10} - N_l}{N_u - N_l} \right) \times C$							
Formula 19	Percentiles in case of Grouped Frequency Distribution: Steps are like median with few modifications.							
	<table border="1"> <thead> <tr> <th>1st Percentile</th> <th>99th Percentile</th> </tr> </thead> <tbody> <tr> <td>Find P_1 class using $\frac{N}{100}$</td> <td>Find P_{99} class using $\frac{99N}{100}$</td> </tr> </tbody> </table>	1 st Percentile	99 th Percentile	Find P_1 class using $\frac{N}{100}$	Find P_{99} class using $\frac{99N}{100}$			
1 st Percentile	99 th Percentile							
Find P_1 class using $\frac{N}{100}$	Find P_{99} class using $\frac{99N}{100}$							



	$P_1 = l_1 + \left(\frac{N - N_1}{N_u - N_1} \right) \times C$ $P_{99} = l_1 + \left(\frac{99N - N_1}{N_u - N_1} \right) \times C$						
Formula 20	<p>Mode in case of discrete observation: observation repeating for maximum no. of times or observation with highest frequency</p> <p>Note: There can be multiple modes also. If all observations are having same frequency, then there is no mode.</p>						
Formula 21	<p>Mode in case of grouped frequency distribution: Find Modal Class (Class with highest frequency) then apply below formula</p> <p>Mode = $l_1 + \frac{f_0 - f_{-1}}{2f_0 - f_{-1} - f_1} \times C$</p> <p>where, l_1 = LCB of modal class, f_0 = frequency of modal class, f_{-1} = frequency of pre-modal class, f_1 = frequency of post modal class, C = class length of modal class</p> <p style="text-align: right;"><i>modal class</i></p> <table style="margin-left: auto; margin-right: auto;"> <tr> <td style="border: 1px solid black; padding: 2px;">20-30</td> <td style="border: 1px solid black; padding: 2px;">30-40</td> <td style="border: 1px solid black; padding: 2px;">40-50</td> </tr> <tr> <td style="text-align: center;">f_{-1}</td> <td style="text-align: center;">f_0</td> <td style="text-align: center;">f_1</td> </tr> </table>	20-30	30-40	40-50	f_{-1}	f_0	f_1
20-30	30-40	40-50					
f_{-1}	f_0	f_1					
Formula 22	<p>Relationship between Mean, Median and Mode in case of Symmetrical Distribution:</p> <p style="text-align: center;">Mean = Median = Mode</p>						
Formula 23	<p>Relationship between Mean, Median and Mode in case of moderately skewed distribution:</p> <p style="text-align: center;">Mean - Mode = 3 (Mean - Median)</p> <p style="text-align: center;">Mode = 3 Median - 2 Mean</p>						
Formula 24	<p>Geometric Mean in case of discrete positive observations:</p> $(x_1 \times x_2 \times x_3 \times \dots \times x_n)^{1/n}$						
Formula 25	<p>Geometric Mean in case of frequency distribution:</p> $(x_1^{f_1} \times x_2^{f_2} \times x_3^{f_3} \times \dots \times x_n^{f_n})^{1/N}$ <p style="text-align: right;">$N = \sum f$</p>						
Formula 26	<p>Harmonic Mean in case of discrete observations:</p> $HM = \frac{n}{\sum (1/x)}$						



Formula 27	Harmonic Mean in case of frequency distribution: $\frac{N}{\sum (f/x)}$	
Formula 28	Combined HM: $H_c = \frac{n_1 + n_2}{\frac{n_1}{H_1} + \frac{n_2}{H_2}}$	
Formula 29	Relationship between AM, GM and HM	
	Situation	Relationship
	When all the observations are identical / same	$AM = GM = HM$
When all the observations are distinct / different	$AM > GM > HM$	
In General	$AM \geq GM \geq HM$	
Formula 30	Range in case of discrete observations: $L - S$ where L = Largest Observation, S = Smallest Observation	
Formula 31	Range in case of Grouped Frequency Distribution: L – S L = UCB of last class interval, S = LCB of first-class interval	
Formula 32	Coefficient of Range $\frac{L - S}{L + S} \times 100$	
Formula 33	Mean Deviation in case of discrete observations (abs dev) $\frac{\sum X - A }{n}$ where A is any appropriate central tendency (as given) $\frac{\sum X}{n}$	
Formula 34	Mean Deviation (in case of grouped frequency distributions) $MD_A = \frac{1}{N} \sum f x - A $ where A is any appropriate central tendency (as given) $N = \sum f$ $\frac{\sum fx}{N}$	



Formula 35	Coefficient of Mean Deviation: $\frac{MD \text{ about } A}{A} \times 100$
Formula 36	Standard Deviation in case of discrete observations: $\sqrt{\frac{\sum (x - \bar{x})^2}{n}}$ or shorter formula $\sqrt{\frac{\sum x^2}{n} - \bar{x}^2}$
Formula 37	Standard Deviation in case of grouped frequency observations $\sigma_x = SD_x = \sqrt{\frac{\sum f(x - \bar{x})^2}{N}}$ or shorter formula $\sigma_x = SD_x = \sqrt{\frac{\sum fx^2}{N} - (\bar{x})^2}$
Formula 38	Coefficient of Variation: $\frac{SD}{Am} \times 100$
Formula 39	If there are only <u>two</u> observations, then SD is half of range $SD = \frac{Range}{2}$
Formula 40	Standard Deviation of first n natural numbers: $\sqrt{\frac{n^2 - 1}{12}}$
Formula 41	Combined SD: $s_c = \sqrt{\frac{n_1 s_1^2 + n_2 s_2^2 + n_1 d_1^2 + n_2 d_2^2}{n_1 + n_2}}$ $\bar{x}_c = \frac{n_1 \bar{x}_1 + n_2 \bar{x}_2}{n_1 + n_2}$ $d_1 = \bar{x}_c - \bar{x}_1 \text{ and } d_2 = \bar{x}_c - \bar{x}_2$
Formula 42	If all the observations are constant, then SD/ MD/ Range is <u>zero</u>
Formula 43	Change of Origin and Scale: No effect of change of origin but affected by change of scale in the magnitude (ignore sign) $SD_y = b SD_x$ Note: same thing will apply to all the measures of dispersion

$$SD_y = SD_x \times |chg \text{ of scale} |$$



Formula 44	Quartile Deviation: $QD = \frac{Q_3 - Q_1}{2}$
Formula 45	Coefficient of Quartile Deviation: $\frac{Q_3 - Q_1}{Q_3 + Q_1} \times 100$
Formula 46	Relationship between SD, MD and QD $4SD = 5MD = 6QD$ OR $SD : MD : QD = 15 : 12 : 10$
Formula 47	Basic Formula of Probability: $P(A) = \frac{\text{No. of favorable events to A}}{\text{Total no. of events}}$
Formula 48	Odds in favour of Event A: $\frac{\text{no. of fav events}}{\text{no. of unfav events}}$
Formula 49	Odds against an Event A: $\frac{\text{no. of unfav events}}{\text{no. of fav events}}$
Formula 50	Number of total outcomes of a random experiment: If an experiment results in <u>p</u> outcomes and if it is repeated <u>q</u> times, then Total <u> </u> of outcomes <u> </u> p^q
Formula 51	Relative Frequency Probability $\frac{\text{no. of times the event occurred during experimental trials}}{\text{total no. of trials}} = \frac{f_A}{n}$
Formula 52	Set Based Probability: $P(A) = \frac{\text{no. of sample points in A}}{\text{no. of sample points in S}} = \frac{n(A)}{n(S)}$ here A is Event Set and S is Sample Space
Formula 53	Addition Theorem 1: In case of two mutually exclusive events A and B $P(A \cup B) = P(A + B) = P(A \text{ or } B) = P(A) + P(B)$
Formula 54	Addition Theorem 2: In case of two or more mutually exclusive events $P(A_1 \cup A_2 \cup A_3 \cup \dots) = P(A_1) + P(A_2) + P(A_3) + \dots$



Formula 55	Addition Theorem 3: For any two events $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
Formula 56	Addition Theorem 4: In case of any three events $P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(B \cap C) - P(A \cap C) + P(A \cap B \cap C)$
Formula 57	Conditional Probability of Event B when Event A is already occurred $P(B/A) = \frac{P(A \cap B)}{P(A)}$ provided $P(A) \neq 0$
Formula 58	Conditional Probability of Event A when Event B is already occurred $P(A/B) = \frac{P(A \cap B)}{P(B)}$ provided $P(B) \neq 0$
Formula 59	Compound Theorem: In case of two dependent events $P(A \cap B) = P(B) \times P(A/B) \text{ or}$ $P(A \cap B) = P(A) \times P(B/A)$
Formula 60	Compound Theorem: In case of two independent events $P(A \cap B) = P(A) \times P(B)$
Formula 61	Expected value of a Probability Distribution: $E(X) = \sum PX$ Also, $E(x) = \mu$ (here μ means mean of probability distribution)
Formula 62	Variance of Probability Distribution: $V(x) = E(x - \mu)^2 = E(x^2) - [E(x)]^2$
Formula 63	Probability Mass Function in case of Binomial Distribution: $f(x) = P(X = x) = {}^n C_x p^x q^{n-x}$



Formula 64	Mean of Binomial Distribution: $\mu = np$
	Variance of Binomial Distribution: $\sigma^2 = npq$
Formula 65	Mode in case of Binomial Distribution:
	Step 1 Calculate $(n+1)p$
	Step 2A If $(n+1)p$ is an integer, there will be two modes: $\mu_0 = (n+1)p$ & $[(n+1)p - 1]$
Step 2B If $(n+1)p$ is a non-integer, there will be only one mode: $\mu_0 =$ largest integer contained in $(n+1)p$	
Formula 66	Probability Mass Function in case of Poisson Distribution: $f(x) = P(X = x) = \frac{e^{-m} \cdot m^x}{x!}$ $e = 2.71828$
Formula 67	Mean of Poisson Distribution: $\mu = m$
	Variance of Poisson Distribution: $\sigma^2 = m$
	SD of Poisson Distribution: $\sigma = \sqrt{m}$
Formula 68	Mode in case of Poisson Distribution:
	If m is an integer there will be two modes: $\mu_0 = m$ & $m-1$
	If m is a non-integer there will be only one mode: largest integer contained in m
Formula 69	Probability Density Function in case of Normal Distribution $f(x) = \frac{1}{\sigma\sqrt{2\pi}} \cdot e^{-\left(\frac{x-\mu}{\sigma}\right)^2 \frac{1}{2}}$
Formula 70	Mean Deviation in case of Normal Distribution: $MD = 0.8 \sigma$
Formula 71	Quartiles in case of Normal Distribution:
	$Q_1 = \mu - 0.675 \sigma$
	$Q_3 = \mu + 0.675 \sigma$



Formula 72	Quartile Deviation in case of Normal Distribution: $QD = 0.675\sigma$	
Formula 73	Points of Inflex of Normal Curve: $\mu - \sigma, \mu + \sigma$	
Formula 74	In case of Normal Distribution, Ratio between QD: MD: SD = 10:12:15	
Formula 75	Conditions of Standard Normal Distribution: Mean = 0, SD = 1 $\mu = 0, \sigma = 1$	
Formula 76	Z Score: $z = \frac{x - \mu}{\sigma}$	
Formula 77	Area under Normal Curve (Popular Intervals)	
	From	To
	μ	$\mu + \sigma$
	$\mu + \sigma$	$\mu + 2\sigma$
	$\mu + 2\sigma$	$\mu + 3\sigma$
	$\mu + 3\sigma$	$+\infty$
	Area under Normal Curve Probability	
		34.135%
		13.59%
		2.14%
		0.135%
Formula 78	For a $p \times q$ bivariate frequency distribution:	
	Number of cells	$p \times q$
	Number of marginal distributions	2
	Number of conditional distributions	$p + q$
Formula 79	Karl Pearson's Product Moment Correlation Coefficient: $r_1 = \frac{cov(x,y)}{SD_x SD_y}$	
Formula 80	Covariance between two variables: $cov(x,y) = \frac{\sum(x-\bar{x})(y-\bar{y})}{n}$ or $\frac{\sum xy}{n} - \bar{x} \cdot \bar{y}$	



<p>Formula 81</p>	<p>Spearman's Rank Correlation Coefficient:</p> $r_R = 1 - \frac{6\sum d^2}{n(n^2-1)}$
<p>Formula 82</p>	<p>here d means difference in ranks of both variables Spearman's Rank Correlation Coefficient (in case of tied values)</p> $r_R = 1 - \frac{6(\sum d^2 + A)}{n(n^2-1)}$ <p>here A is adjustment value</p> $A = \frac{\sum(t^3 - t)}{12}$ <p>where t = tie length (calculate t value for each of the ties)</p>
<p>Formula 83</p>	<p>Coefficient of Concurrent Deviations</p> $r_c = \frac{+}{\sqrt{\frac{+}{m}(\frac{QC-m}{m})}}$ <p>n = total pair</p> <p>where c is number of concurrent deviations (same direction) m is number of pairs compared (equals to n-1)</p>
<p>Formula 84</p>	<p>Regression Coefficients:</p> <p>Y on X:</p> $b_{YX} = r \frac{SD_y}{SD_x}$ $b_{YX} = \frac{cov(x,y)}{var\ of\ x}$ <p>X on Y:</p> $b_{XY} = r \frac{SD_x}{SD_y}$ $b_{XY} = \frac{cov(x,y)}{var\ of\ y}$
<p>Formula 85</p>	<p>Correlation Coefficient is the GM of regression coefficients:</p> $r_{xy} = \pm \sqrt{b_{YX} \times b_{XY}}$ <p>Note: r_{xy}, b_{xy}, b_{yx} all will have same sign</p>



Formula 86	<p>Change of Origin/ Scale for Regression Coefficients: Origin no impact, Scale impact of both magnitude and sign.</p> <p>$x \rightarrow u$ $b_{vu} = b_{yx} \times \frac{\text{change of scale of } y}{\text{change of scale of } x}$ b_{vy} b_{yx}</p> <p>$y \rightarrow v$ $b_{uv} = b_{xy} \times \frac{\text{change of scale of } x}{\text{change of scale of } y}$ b_{uv} b_{xy}</p>
Formula 87	<p>Two regression lines (if not identical) will intersect at the point (\bar{x}, \bar{y})</p>
Formula 88	<p>Coefficient of Determination/ Explained Variance/ Accounted Variance: r^2</p>
Formula 89	<p>Coefficient of Non-determination/ Un-explained Variance/ Un-accounted Variance: $1 - r^2$</p>
Formula 90	<p>Price Relatives: $\frac{P_n}{P_0}$, Quantity Relatives: $\frac{Q_n}{Q_0}$, Value Relatives: $\frac{V_n}{V_0}$</p>
Formula 91	<p>Simple Aggregative Index: $\frac{\sum P_n}{\sum P_0} \times 100$</p>
Formula 92	<p>Simple Average of <u>Relatives</u> – Method Index: $\frac{\sum \frac{P_n}{P_0}}{n}$</p>
Formula 93	<p>Laspeyres Index (weight – base year quantity weight) q_0</p> <p>$\frac{\sum P_n q_0}{\sum P_0 q_0} \times 100$</p>
Formula 94	<p>Paasche's Index (weight – current year quantity weight) q_n</p> <p>$\frac{\sum P_n q_n}{\sum P_0 q_n} \times 100$</p>



Formula 95	Marshall-Edgeworth Index (weight – sum of both current and base quantity) $\frac{\sum P_n (Q_0 + Q_n)}{\sum P_0 (Q_0 + Q_n)} \times 100$
Formula 96	Fisher's Ideal Index: GM of Laspeyres Index and Paasche's Index $\sqrt{L \times P}$
Formula 97	Bowley's Index: AM of Laspeyres Index and Paasche's Index $\frac{L + P}{2}$

About CA. Pranav Popat Sir

- He is a Chartered Accountant (Inter and Final Both Groups in First Attempt) with 7+ years of experience.
- He is an Educator by Passion and his Choice (Dil Se ♥)
- He teaches subjects of Maths, LR and Stats (Paper 3) at CA Foundation Level and Cost & Management Accounting (Paper 3) at CA Intermediate Level.

Hope this formula book helps you in revising all formulas and become helpful to you during exam time, I made this with my whole heart, make best use of it and I just want one thing in return – share these notes to every student who really needs this.

Wishing you ALL THE BEST for upcoming examinations, see you soon in Inter Costing!!!

Ab mushkil nahi kuch bhi, nahi kuch bhi!!!

With Lots of Love

CA. Pranav Popat (P² SIR)



CA INTERMEDIATE MAY 25



Early Bird Batch



CA Deepika Rathi
Corp. Law & Auditing

Starting
8th July



CA Indresh Gandhi
Corp. Law & Auditing



CA Vivek Gaba
Taxation



CA Pranav Popat
Costing



CA Mohnish Vora
FM/SM



CA Tejas Suchak
Adv. Accounting