

Inequalities & Equations



1. The standard format of a linear equation is : $ax + by + c = 0$
 where $a, b \neq 0$ at a time.


OR $y = mx + c$ where $m = \text{slope of the line}$

2.

Linear Equation	a	b	c
$5x + 13y + 8 = 0$	5	13	8
$133x - 18y + 2k - 18 = 0$	133	-18	$2k - 18$
$(2p+3)x - (18ky) + 339 = 93$	$(2p+3)$	$-18k$	246
$2x + 3y = 88$	2	3	-88
$17kx + 5x - 3y = 18k - 23$ $(17k+5)x - 3y - 18k + 23 = 0$	$(17k+5)$	-3	$-18k + 23$
$2x = 83$ i.e. $2x + 0y - 83 = 0$	2	0	-83
$8y = -2k - 99$ i.e. $0x + 8y + 2k + 99 = 0$	0	8	$2k + 99$
$kx - 13y - 25y - 9y + t = 0$ i.e. $kx - 47y + t = 0$	k	-47	t
$5x + 13y = 27x - 40y + 88$ i.e. $-22x + 53y - 88 = 0$	-22	53	-88
$2px - 13y + 8kx - 11py - 33$ $= 12mx - 18py - 11$ i.e. $(2p+8k-12m)x + (-13-11p+18p)y - 22 = 0$	$(2p+8k-12m)$	$(-13-11p+18p)$	-22



3.



4.

Points	Location	Equation / Inequalities
(+, +)	1st quadrant	$x > 0, y > 0$
(-, +)	2nd quadrant	$x < 0, y > 0$
(-, -)	3rd quadrant	$x < 0, y < 0$
(+, -)	4th quadrant	$x > 0, y < 0$
(±, 0)	x-axis	$y = 0$
(0, ±)	y-axis	$x = 0$
(0, 0)	origin	$x, y = 0$



- Equation of X-axis is $y = 0$
- Equation of Y-axis is $x = 0$
- $(0, 0)$ represents origin = Point of intersection of X, Y -axis
- If x - co-ordinate of a point is 0, then that point is on : **Y-axis**
 examples : $(0, 30), (0, 20), (0, -80), (0, -25), (0, \frac{2}{3})$ are on Y-axis
- If y - co-ordinate of a point is 0, then that point is on : **X-axis**
 examples : $(30, 0), (20, 0), (45, 0), (-25, 0), (-28.80, 0)$ these points are on X-axis

5. Find points satisfying the linear equation $2x + 3y = 300$

If I put $x = 150, y = 0$ then $2x + 3y = 300$ is satisfied

therefore, $(150, 0)$ is one of the point satisfying the equation $2x + 3y = 300$

Other points : $(0, 100), (10, \frac{280}{3}), (20, \frac{260}{3}), (15, 90), (300, -100), (60, 60), (75, 50), (40, \frac{220}{3}), (-19, \frac{338}{3}), \dots$

Such infinite points can satisfy this linear equation.


6. Find points satisfying the linear equation $x + y = 50$

➔ $(0, 50), (50, 0), (30, 20), (20, 30), (10, 40), (25, 25), (60, -10), (-20, 70)$
 $(-100, 150), (250, -200), (1.50, 48.50), (2.85, 47.15), (1, 49), (15, 35), (45, 5), \dots$

such infinite points can satisfy this equation.

7. Find points satisfying the linear equation $2x + y = 100$ and plot those points on graph paper?

➔ Points satisfying $2x + y = 100$ are : $(0, 100), (50, 0), (25, 50), (10, 80), (20, 60), (30, 40), (40, 20), (60, -20), (70, -40), (35, 30), (15, 70), (5, 90), (-10, 120), (-20, 140), (-30, 160) \dots$



Line is a set / collection of infinite points satisfying the given. Linear equation

Graphical presentation of a linear equation is known as Line.

8. How to draw a line on a graph paper if equation of the line is given?

- ➔
- ① Find atleast 2 points satisfying the given Linear Equation
 - ② plot those points on Graph paper
 - ③ Draw a straight line passing through all the points


Graphical representation of Linear equation is line

Line is a set/collection of infinite points satisfying the given linear equation.

9. Draw the line $5x + 6y = 240$ on graph paper



points satisfying Eqⁿ $5x + 6y = 240$: $(0, 40), (48, 0)$




10. Draw the line $15x - 2y = 300$ on graph paper



points satisfying the Eqⁿ $15x - 2y = 300$

: $(20, 0), (0, -150)$



11. Draw the line $y = 30$ on Graph paper.




points satisfying eqⁿ $y = 30$ are $(0,30), (10,30), (20,30), (30,30), (40,30), (-10,30), (-20,30), (-30,30), (-40,30), (-50,30), (-60,30), (-70,30), (-80,30), (-90,30), (-100,30)$
 $(0x + y = 30)$



▪ If Eqⁿ of the line is $y = \text{constant}$ then that line is || to $X\text{-axis}$

▪ If Eqⁿ of the line is $x = \text{constant}$ then that line is || to $Y\text{-axis}$

12.



Eqⁿ of X-axis : $y = 0$


Eqⁿ of Y-axis : $x = 0$



13. Draw the lines $(x + y = 50)$ & $(2x + y = 60)$ on graph paper and Find point of intersection of these 2 lines.

→ $x + y = 50 \implies (0, 50), (50, 0)$

$2x + y = 60 \implies (0, 60), (30, 0)$



on solving 2 linear Eqns simultaneously if we get $x = p, y = q$ then (p, q) is the point of intersection of 2 lines

$$\begin{array}{r} 2x + y = 60 \text{ ----- ①} \\ - \quad x + y = 50 \text{ ----- ②} \\ \hline x = 10 \end{array}$$

$$\begin{array}{l} x + y = 50 \\ 10 + y = 50 \\ y = 40 \end{array}$$


$(10, 40)$ is the point of intersection.



14. Draw the lines $2x + 3y = 60$ and $5x + 2y = 95$ on graph paper and Find point of intersection.

$$2x + 3y = 60 : (30, 0), (0, 20)$$

$$5x + 2y = 95 : (25, -15), (35, -40)$$



To get the point of intersection, Let's solve 2 linear equations simultaneously : $2x + 3y = 60$ & $5x + 2y = 95$

Let's multiply eqⁿ ① by 2 on both sides, eqⁿ ② by 3 on both sides

$$\begin{array}{r} 4x + 6y = 120 \\ -15x + 6y = 285 \\ \hline -11x = -165 \\ \boxed{x = 15} \end{array}$$

on solving 2 linear eqⁿs simultaneously if we get $x = m$, $y = n$ then (m, n) is the point of intersection of 2 Lines. If we put $x = m$, & $y = n$ the Both eqⁿs must be satisfied

Let's put $x = 15$ in one of the equation,

$$2x + 3y = 60$$

$$2(15) + 3y = 60$$

$$3y = 30$$

$$\boxed{y = 10}$$

$\therefore (15, 10)$ is the point of intersection of point.

15. Find point of intersection of $3x + 5y = 90$ & $2x + 3y = 60$



$$\begin{array}{r}
 9x + 15y = 270 \\
 10x + 15y = 300 \\
 \hline
 -x = -30 \\
 x = 30
 \end{array}$$

$$3x + 5y = 90$$

$$3(30) + 5y = 90$$

$$5y = 0$$

$$y = 0$$

$\therefore (30, 0)$ is the point of intersection.

16. Find point of intersection of lines $3x + 5y = 100$ & $5x + 3y = 150$



$$\begin{array}{r}
 15x + 25y = 500 \\
 15x + 9y = 450 \\
 \hline
 16y = 50
 \end{array}$$

$$16y = 50$$

$$y = 3.125$$

$$3x + 5(3.125) = 100$$

$$x = 28.125$$

$$\left(\frac{225}{8}, \frac{25}{8}\right)$$



$\therefore (28.125, 3.125)$ is the point of intersection of 2 lines.

17. Point of intersection of $7x - 3y = 20$ & $5x + 13y = 200$ lie in

_____ Quadrant.



$$\begin{array}{r}
 35x - 15y = 100 \\
 35x + 91y = 1400 \\
 \hline
 -106y = -1300
 \end{array}$$

$$-106y = -1300$$

$$y = 12.264151$$

$$7x - 3(12.264151) = 20$$

$$x = 8.11321$$

$(8.11321, 12.264151)$ is the point of intersection. (i.e. 1st quadrant)



18. Point of intersection of lines $5x + 2y = 90$, $10x + 9y = 180$ lie in _____ Quadrant

a. 1st

b. 2nd

c. 3rd

~~d. None of these~~

$$\begin{array}{r} 10x + 4y = 180 \\ 10x + 9y = 180 \\ \hline -5y = 0 \\ y = 0 \end{array}$$

$\therefore (18, 0)$ is the point of intersection.
i.e. point of intersection is on x-axis

$$\begin{aligned} \therefore 5x + 2(0) &= 90 \\ x &= 18 \end{aligned}$$

19. Find point of intersection of $8x - y = 90$ & $3x - 7y = 190$



$$\begin{array}{r} 56x - 7y = 630 \\ 3x - 7y = 190 \\ \hline 53x = 440 \end{array}$$

$$x = 8.301887$$

$$8(8.301887) - y = 90$$

$$\therefore y = -23.58491$$

point of intersection is $(8.30, -23.58)$ which is in 4th quadrant


20. The point $(8k, -19)$ lie on the line $5x - 13y = 88$. Find value of k.



$$5(8k) - 13(-19) = 88$$

$$40k + 247 = 88$$

$$k = -3.975$$



21. The point $(-\frac{k}{3}, 35)$ lie on the line $10x - 55y = 230$. Find value of k.

As point $(-\frac{k}{3}, 35)$ lie on the line

$$10x - 55y = 230$$


\therefore If we put $x = -\frac{k}{3}$ & $y = 35$

then Given eqⁿ must be satisfied

$$\therefore 10x - 55y = 230$$

$$10\left(-\frac{k}{3}\right) - 55(35) = 230$$

$$\therefore -\frac{10k}{3} = 2155 \quad \therefore k = -646.50$$



22. Find point of intersection of lines $2x + 3y = 800$ and


$$8x + 12y = 1000$$



$$2x + 3y = 800$$

$$8x + 12y = 3,200$$

$$8x + 12y = 1000$$



Slope of the line $ax + by + c = 0$ is $-\frac{a}{b}$

Slope of the line $2x + 3y = 800$ is $-\frac{2}{3}$

Slope of the line $8x + 12y = 1000$ is $-\frac{8}{12} = -\frac{2}{3}$

As the slope of 1st line = slope of 2nd line, Lines are || to each other.

$$-\frac{2}{3} = -\frac{2}{3}$$


If m_1 is slope of one line & m_2 is slope of other line then lines are said to be || to each other if $m_1 = m_2$

23. Draw the lines $3x - 2y = 60$ & $9x - 6y = 300$ on graph paper.

Find point of intersection.

➔ $3x - 2y = 60$ ➔ $(20, 0), (0, -30)$

$9x - 6y = 300$ ➔ $(0, -50), (30, -5)$



There is no point of intersection as these lines are || to each other

Slope of 1st line = Slope of 2nd line = $\frac{3}{2}$
 $-\frac{3}{-2} = -\frac{9}{-6} = \frac{3}{2}$

24. The lines $5x + 11y = 22$ and $8kx - 55y = -980$ are || to each other.

Find value of k.

➔ As these 2 lines are || to each other
 slope of 1st line = slope of 2nd line

$$\frac{-5}{11} = \frac{-8k}{-55} \quad \therefore \frac{-5}{11} = \frac{8k}{55}$$

$$\therefore 88k = -275 \therefore k = -3.125$$

25. The lines $5x + 13y = 80$ and $8mx - 22y = 810$ are || to each other.

Find value of m.

➔ As these 2 lines are || to each other

Inequalities & Equations



Slope of 1st line = slope of 2nd line

$$-\frac{5}{13} = \frac{-8m}{-22}$$

$$\therefore 104m = -110$$

$$\frac{-5}{13} = \frac{8m}{22}$$

$$m = \frac{-110}{104} = \frac{-55}{52} = -1.05769$$

26.

Eq ⁿ of the line	Slope of the line
$ax + by - c = 0$	$-a/b$
$3x + 5y + 30 = 0$	$-3/5$
$3x + 5y - 1000 = 0$	$-3/5$
$5x - 13y = 88$	$-a/b = -5/-13 = 5/13$
$8kx - 33py = 8k - p$	$8k/33p$
$29x - 33y = 5x - 88$	$24x - 33y + 88 = 0 \therefore \text{Slope} = \frac{24}{33} = \frac{8}{11}$
$24x - 33y = -88$	
$13x - 2y = 88x - 130y - y + 2x$ $-8p + 63$	$\text{Slope} = \frac{77}{129}$
i.e. $-77x + 129y = -8p + 63$	
$31x - 2y = 8kx - 55y + 11$	$\text{slope} = \frac{-(31-8k)}{53} = \frac{8k-31}{53}$
i.e. $(31-8k)x + 53y = 11$	
$x = 35$	$\text{slope} = \frac{-1}{0} = \text{Not defined}$
i.e. $x + 0y - 35 = 0$	$= \text{undefined}$
$2x = 101$	$\text{slope} = \frac{-2}{0} = \text{undefined}$
$2x + 0y - 101 = 0$	
$5y = 33$	$\text{slope} = \frac{-0}{5} = 0 = \text{Zero}$
$0x + 5y = 33$	
$0x + y = 33$	$\text{slope} = \frac{-0}{1} = 0 = \text{Zero}$
$x = 500$	i.e. $x + 0y = 500 \therefore \text{slope} = -1/0 = \text{Not defined}$
$px + qy + r = 0$	$-p/q$
$33x + py = r$	$-33/p$

27. Find slope of line $x = 155$ i.e. $x + 0y = 155$

Slope = $-1/0 = \text{Not defined}$

28. Find slope of line $y = 30$ i.e. $0x + y = 30$

Slope = $-0/1 = \text{zero} = 0$

Slope of X-axis and all the lines || to X-axis is : Zero

Slope of Y-axis and all the lines || to Y-axis is : undefined or Not defined

A line	Slope	Equation
X - Axis	Zero	$y = 0$
Y - Axis	Not defined	$x = 0$
to X - Axis	Zero	$y = \text{constant}$
to Y - Axis	Not defined	$x = \text{constant}$

29. Standard format of a linear equation is,

$$ax + by + c = 0$$

$$by = -ax + c$$

dividing by b on both sides

$$\frac{by}{b} = \frac{-ax + c}{b}$$

$$y = \left(\frac{-a}{b}\right)x + \text{constant}$$

$$y = mx + c$$

where, m = slope of the line.

$$y = 8x + 13 \implies m = \text{slope} = 8$$

$$-8x + y - 13 = 0 \implies \text{slope} = -a/b = 8$$

slope of the line

$$3x + 5y = 88 \text{ is } -3/5$$



$$3x + 5y = 88$$

$$5y = 88 - 3x$$

$$5y = -3x + 88$$

$$y = \left(-\frac{3}{5}\right)x + \left(\frac{88}{5}\right)$$

comparing this with

$$y = mx + c$$

$$m = -3/5 = \text{slope of line}$$

30. Find slope of the line $3x + 5y = 88$

$$3x + 5y - 88 = 0$$

comparing this with $ax + by + c = 0$

$$a = 3, b = 5$$

$$\therefore \text{slope of the line} = -a/b$$

$$= -3/5$$

$$3x + 5y = 88$$

$$5y = 88 - 3x$$

dividing by 5 on both sides

$$y = -\frac{3}{5}x + \frac{88}{5}$$

comparing this with $y = mx + c$

$$m = -\frac{3}{5} = \text{slope of the line}$$

31. Find any 2 points satisfying the equation $7x - 3y = 100$

→ 2 points satisfying the equation $7x - 3y = 100$ are : $(100, 200), (10, -10)$

32. Find eqⁿ of the line passing through points $(100, 200)$ & $(10, -10)$



$$\begin{matrix} (100, 200) & & (10, -10) \\ \downarrow & & \downarrow \\ (x_1, y_1) & & (x_2, y_2) \end{matrix}$$

$$\frac{y_2 - y_1}{y - y_1} = \frac{x_2 - x_1}{x - x_1} \quad \dots \text{Eq}^n \text{ of line passing through } (x_1, y_1) \text{ \& } (x_2, y_2)$$

$$\frac{-10 - 200}{y - 200} = \frac{10 - 100}{x - 100}$$

$$\frac{-210}{y - 200} = \frac{-90}{x - 100}$$

$$-210(x - 100) = -90(y - 200)$$

$$-210x + 21000 = -90y + 18000$$

$$-210x + 90y = -3000$$

$$210x - 90y = 3000$$

$$7x - 3y = 100$$

33. Find equation of the line passing through points $(x_1, y_1), (x_2, y_2)$

→ Let $y = mx + c$ be the eqⁿ of the line passing through point


(x_1, y_1) & (x_2, y_2)

$y = mx + c$ ①

As point (x_1, y_1) is on the line $y = mx + c$

$y_1 = mx_1 + c$ ②

Similarly, $y_2 = mx_2 + c$ ③



eqⁿ ① - eqⁿ ②

$y - y_1 = mx + c - mx_1 - c$

$y - y_1 = m(x - x_1)$

∴ $m = \frac{y - y_1}{x - x_1}$ ④

eqⁿ ③ - eqⁿ ②

$y_2 - y_1 = mx_2 + c - mx_1 - c$

$y_2 - y_1 = m(x_2 - x_1)$

∴ $m = \frac{y_2 - y_1}{x_2 - x_1}$ ⑤

From eqⁿ ④ & eqⁿ ⑤

$\frac{y_2 - y_1}{x_2 - x_1} = \frac{y - y_1}{x - x_1}$

$\left(\frac{y_2 - y_1}{y - y_1}\right) = \left(\frac{x_2 - x_1}{x - x_1}\right) \dots \dots \dots \rightarrow$

This is eqⁿ of the line passing through points $(x_1, y_1), (x_2, y_2)$

34. Find Eqⁿ of the line passing through point $(p, q), (m, n)$

→ Eqⁿ of line passing through points (x_1, y_1) & (x_2, y_2)


is $\left(\frac{y_2 - y_1}{y - y_1}\right) = \left(\frac{x_2 - x_1}{x - x_1}\right)$

∴ Eqⁿ of line passing through points $(p, q), (m, n)$

is $\left(\frac{n - q}{y - q}\right) = \left(\frac{m - p}{x - p}\right)$



35. Find slope of the line $y = \frac{-8}{5}x + 33$



$$y = \frac{-8}{5}x + 33$$

Comparing this with $y = mx + c$

$$m = \frac{-8}{5} = \text{Slope of the line.}$$

$$y = -\frac{11}{6}x - 99 \text{ ----- slope} = -\frac{11}{6}$$

$$6y = -11x - 594$$

$$11x + 6y + 594 = 0 \text{ ----- slope} = -\frac{a}{b} = -\frac{11}{6}$$

$$y = \frac{-8}{5}x + 33$$

$$y - 33 = \frac{-8}{5}x$$

$$5y - 165 = -8x$$

$$8x + 5y = 165$$

$$8x + 5y - 165 = 0$$

Comparing this with $ax + by + c = 0$

$$a = 8, b = 5$$

$$\text{Slope} = \frac{-a}{b} = \frac{-8}{5}$$

36. Slope of the line $kx + 15y = 2x - 93$ is $\frac{-8}{11}$. Find k.



$$kx + 15y - 2x + 93 = 0$$

$$(k-2)x + 15y + 93 = 0$$

comparing this with $ax + by + c = 0$

$$a = (k-2), b = 15, c = 93$$

$$\text{slope} = \frac{-a}{b} = \frac{-(k-2)}{15} = \frac{-8}{11}$$

$$11(k-2) = 120$$

$$11k - 22 = 120$$

$$11k = 142$$

$$k = \left(\frac{142}{11}\right) = 12.90909090$$

37. Slope of the line $19x - 33y + 2ky = 8x - 930$ is $\frac{1}{8}$. Find k.



$$\rightarrow 19x - 33y + 2ky - 8x + 930 = 0$$

$$11x + (2k - 33)y + 930 = 0$$

comparing this with $ax + by + c = 0$

$$a = 11, \quad b = (2k - 33)$$

$$\text{slope} = \frac{-a}{b} = \frac{-11}{2k - 33} = \frac{11}{8}$$

$$\therefore 22k = 275$$

$$k = \left(\frac{275}{22}\right)$$

$$22k - 363 = -88$$

$$k = 12.50$$

38. Find Eqⁿ of the line passing through points (8, -12), (18, 33)



$$\downarrow$$

$$(x_1, y_1), (x_2, y_2)$$

Eqⁿ of the line passing through (x_1, y_1) & (x_2, y_2) is

$$\frac{y_2 - y_1}{y - y_1} = \frac{x_2 - x_1}{x - x_1}$$

$$45x - 360 = 10y + 120$$

$$45x - 10y = 480$$


$$\frac{33 - (-12)}{y - (-12)} = \frac{18 - 8}{x - 8}$$

$$45x - 10y - 480 = 0$$

$$9x - 2y - 96 = 0$$

$$45(x - 8) = 10(y + 12)$$

39. Find Eqⁿ of the line passing through points (-30, -20), (-1.50, 80)



$$\frac{y_2 - y_1}{y - y_1} = \frac{x_2 - x_1}{x - x_1}$$

$$(x_1, y_1) \quad (x_2, y_2)$$

$$100x + 3000 = 28.50y + 570$$

$$\frac{80 - (-20)}{y - (-20)} = \frac{-1.50 - (-30)}{x - (-30)}$$

$$100x - 28.50y + 2430 = 0$$

$$1000x - 285y + 24300 = 0$$

$$100(x + 30) = 28.50(y + 20)$$

$$200x - 57y + 4860 = 0$$

40. Find Eqⁿ of the line passing through points (2, -5), (-11, 20).

Also find slope of that line

$$\frac{y_2 - y_1}{y - y_1} = \frac{x_2 - x_1}{x - x_1}$$

$$\frac{20 - (-5)}{y - (-5)} = \frac{-11 - 2}{x - 2}$$

$$25x - 50 = -13y - 65$$

$$25x + 13y + 15 = 0$$

Slope of the line = $-\frac{25}{13}$

Slope of the line passing through points (x_1, y_1) & (x_2, y_2) is

$$\left(\frac{y_2 - y_1}{x_2 - x_1} \right)$$

$$\text{Slope of the line} = \frac{20 - (-5)}{-11 - 2} = \frac{25}{-13}$$

Eqⁿ of the line:

$$25x + 13y = 25(2) + 13(-5)$$

$$25x + 13y + 15 = 0$$

41. Find Slope, Eqⁿ of the line passing through points (20, 28), (30, 85).



$$\text{Slope} = \frac{85 - 28}{30 - 20} = \frac{y_2 - y_1}{x_2 - x_1}$$

$$= \left(\frac{57}{10} \right)$$

Eqⁿ of the line

$$57x - 10y = 57(20) - 10(28)$$

$$57x - 10y - 860 = 0$$

OR

$$\frac{y_2 - y_1}{y - y_1} = \frac{x_2 - x_1}{x - x_1}$$

$$\frac{85 - 28}{y - 28} = \frac{30 - 20}{x - 20}$$

$$57(x - 20) = 10(y - 28)$$

$$57x - 1140 = 10y - 280$$

$$57x - 10y = 860$$

$$\therefore \text{slope} = \left(\frac{57}{10} \right)$$

42. Find Slope, Eqⁿ of the line passing through points

(1.50, 18.50), (-27, 35).



$$\text{Slope} = \frac{35 - 18.50}{-27 - 1.50} = \frac{16.50}{-28.50} = \frac{165}{-285} = \frac{33}{-57} = \frac{11}{-19} = \frac{-11}{19}$$

Equation of the line : $11x + 19y = 11(-27) + 19(35)$

$$\therefore 11x + 19y = 368$$

OR $-11x - 19y = -11(-27) - 19(35) = -368$

43. Find Slope of the line passing through points (a, b) & (c, d)

a. $\frac{(d-b)}{(c-a)}$

b. $\frac{(b-d)}{(a-c)}$

~~c. Both~~

d. None

44. Slope of the line passing through (2k, 19) & (50, -8) is $-\frac{16}{3}$

Find k.



Slope of the line passing through points (2k, 19) & (50, -8) = $\frac{(-8-19)}{(50-2k)} = -\frac{16}{3}$

$$\frac{+27}{50-2k} = \frac{+16}{3}$$

$$16(50-2k) = 81$$

$$800 - 32k = 81$$

$$719 = 32k$$

$$\therefore k = \left(\frac{719}{32}\right) = 22.46875$$

45. The line $8x - 3y = 20$ & $7kx + 55y = 250$ have no solution. Find k



As these 2 lines have no solution, we can say that these 2 lines have same slope.

Means point of intersection

Slope of 1st line = Slope of 2nd line

$$\frac{8}{3} = \frac{-7k}{55}$$

$$-21k = 440$$

$$k = \left(\frac{-440}{21}\right) = -20.9523809523$$

46. The lines $5x + 11y = 29$ & $kx + 33y = 810$ have unique solution then

a. $k = 15$

~~b. $k \neq 15$~~

c. $k = 0$

d. Wrong qs.



As these 2 lines have unique solution, It means they have a point of intersection.

Slope of 1st line \neq slope of 2nd line

$$\frac{-5}{11} \neq \frac{-k}{33}$$

$$-11k \neq -165$$

$$k \neq 15$$

47. If m_1 is slope of one line & m_2 is slope of other line then

Lines are said to be

|| to each other
when

$$m_1 = m_2$$

Oblique
when

$$m_1 \neq m_2$$

\perp to each other
when

$$m_1 = \frac{1}{m_2}$$

$$m_1 \cdot m_2 = -1$$

48. $3x - 19y = 50$ & $2kx + 51y = 200$ are \perp to each other.

Find value of k.



As these 2 lines are \perp to each other,

slope of 1st line \times slope of 2nd line = -1

$$\frac{3}{19} \times \frac{-2k}{51} = -1$$

$$\frac{-6k}{969} = -1$$

$$6k = 969 \quad \therefore k = 161.50$$

49.	Slope of the line	Slope of its line	Slope of its ⊥ line
	$\frac{3}{5}$	$3/5$	$-5/3$
	$-\frac{8}{9}$	$-8/9$	$9/8$
	8	8	$-1/8$
	-11	-11	$1/11$
	$\frac{33}{8}$	$33/8$	$-8/33$
	$-\frac{p}{q}$	$-p/q$	q/p
	$\frac{p-q}{r}$	$\frac{p-q}{r}$	$\frac{r}{q-p}$
	0	0	undefined
	Not defined	Not defined	0
	$\frac{3}{91}$	$3/91$	$-91/3$

50. The lines $18x - my = 20$ & $51x - 28y = 290$ are ⊥ to each other.

Find the value of m.



As these 2 lines are ⊥ to each other,

slope of 1st line x slope of 2nd line = -1

$$\frac{18}{m} \times \frac{51}{28} = -1$$

$$918 = -28m$$

$$m = -\frac{918}{28} = -459/14$$

$$m = -32.7857142857$$

51. Draw the line $3x + 5y = 150$ & $5x - 3y = 30$ on graph paper


$\rightarrow 3x + 5y = 150 \rightarrow (50, 0) (0, 30)$

$5x - 3y = 30 \rightarrow (0, -10), (30, 40)$

product of slopes of 2 lines

$= \frac{5}{3} \times -\frac{3}{5}$

$= -1$



slope of this line = $\frac{5}{3}$

These lines are \perp to each other as $m_1 \times m_2 = -1$

slope of this line = $-\frac{3}{5}$

$15x + 25y = 750$

$-15x + 9y = -90$

$34y = 660$

$y = 19.41176$

$\therefore 3x + 5(19.4117647058) = 150$

$x = 17.6471$

52. The lines $8x - 3ky + 21y = 33$ & $15x - 28y = 233$ are \perp to each other. Find k.

$\rightarrow 8x - 3ky + 21y = 33$

$8x + (-3k + 21)y = 33$

$15x - 28y = 233$

slope of the line = $-\frac{8}{-3k+21} = \left(\frac{8}{3k-21}\right)$

slope of the line = $\frac{15}{28}$

As these 2 lines are \perp to each other

$\therefore \frac{8}{3k-21} \times \frac{15}{28} = -1$

$\therefore 120 = -84k + 588$

$k = \frac{-468}{-84} = \frac{117}{21}$

$120 = -28(3k-21)$

$k = 39/7$

Question

$$5x + 2y = 93 \text{ and } 7kx + 12x - 13y - 98y = 200$$

are \perp to each other. Find k .



$$5x + 2y = 93 \text{ \& } (7k+12)x - 111y = 200 \text{ are } \perp \text{ to each other}$$

$$\therefore \left(\begin{array}{l} \text{slope of 1st} \\ \text{line} \end{array} \times \begin{array}{l} \text{slope of} \\ \text{2nd line} \end{array} \right) = -1 \quad \therefore k = \left(\frac{162}{35} \right)$$

$$\frac{+5}{2} \times \frac{7k+12}{111} = +1$$

$$k = 4.628571$$

$$35k + 60 = 222$$

53. Find Eqⁿ of line passing through point (8, 20) having slope (-0.60)



$$\text{Slope} = (-0.60 / 1)$$

$$\text{Slope} = -0.60$$

Eqⁿ of the line

$$= -3/5$$

$$0.60x + y = 0.60(8) + 20$$

Eqⁿ of the line

$$0.60x + y = 24.80$$

$$3x + 5y = 3(8) + 5(20)$$

$$6x + 10y = 248$$

$$3x + 5y = 124$$

$$3x + 5y = 124$$

54. Find Eqⁿ of the line passing through point (8, 10) having slope of 0.70.



$$\text{slope of the line} = 0.70 = \frac{7}{10}$$

$$\therefore \text{Eqⁿ of the line is: } 7x - 10y = 7(8) - 10(10)$$

$$7x - 10y = -44$$

(OR)


$$-7x + 10y = -7(8) + 10(10)$$

$$-7x + 10y = 44$$

$$7x - 10y = -44$$

$$7x - 10y + 44 = 0$$

55. Find eqⁿ of line passing through point (12, 20) and \perp to line joining (21, 6), (18, -9)



slope of $L_1 = -\frac{1}{5}$


Eqⁿ of L_1

$$x + 5y = 12 + 5(20)$$

$$x + 5y = 112$$

56. Find eqⁿ of line passing through point (-16, 12) and \perp to

$$8x - 33y = 200$$



slope of $L_1 = -\frac{33}{8}$


Eqⁿ of L_1

$$33x + 8y = 33(-16) + 8(12)$$

$$33x + 8y = -432$$

$$33x + 8y + 432 = 0$$

57. Find Eqⁿ of the line passing through point (8, -25) & \parallel to line joining (18, 10), (26, -12)



slope of $L_1 = -\frac{11}{4}$

∴ Eqⁿ of L_1

$$11x + 4y = 11(8) + 4(-25)$$

$$11x + 4y = -12$$

$$11x + 4y + 12 = 0$$

58. Find Eqⁿ of line having slope $\frac{8}{5}$ & passing through points (20,16)

→

Eqⁿ of Line : $8x - 5y = 80$
OR
 $-8x + 5y = -80$

59. Find eqⁿ of line passing through point (0.50, 8.75) and \perp to $17x - 20y = 88$

→

\therefore Eqⁿ of L_1 : $20x + 17y = 20(0.50) + 17(8.75)$
 $20x + 17y = 158.75$
 $80x + 68y = 635$

60. If slope of line is zero then that line can be

- a. X-Axis b. \parallel to X-Axis c. \perp to Y-Axis d. All of these

61. If slope of line is Not Defined then that line can be

- a. Y-Axis b. \parallel to Y-Axis c. \perp to X-Axis d. All of these

62. The line $x = 25/2$ is _____

- a. \parallel to Y-Axis b. \perp to X-Axis c. Both d. None




63. Find Eqⁿ of line passing through points (8, 5), (9, 5)



$$0x + y = 5$$

$$y = 5$$




64. Find Eqⁿ of line passing through points (6, 0), (19, 0)



$$y = 0$$

65. Find Eqⁿ of line passing through points (0, 18), (18, 0)



$$x + y = 18$$

$$\text{slope} = \frac{0-18}{18-0} = \frac{-18}{18} = -\frac{1}{1}$$

66. Find Eqⁿ of line passing through points (0, 19), (5, 19)



$$y = 19$$

67. Slope of line passing through points $(\frac{8}{3}, \frac{7}{5}), (\frac{2k}{7}, \frac{19}{3})$ is $\frac{5}{11}$. Find k.



Slope of the line passing

through points $(\frac{8}{3}, \frac{7}{5})$ & $(\frac{2k}{7}, \frac{19}{3})$

$$= \frac{\frac{19}{3} - \frac{7}{5}}{\frac{2k}{7} - \frac{8}{3}}$$

$$\frac{5}{11} = \frac{\frac{95-21}{15}}{\frac{6k-56}{21}}$$

$$\therefore \frac{5}{11} = \frac{74}{15} \times \frac{21}{6k-56}$$

$$6k-56 = \frac{74}{15} \times \frac{21}{1} \times \frac{11}{5}$$

$$k = 47.32$$

68. The lines $3kx - 22y = 80$ & $90x - 47y = 285$ are \perp to each other.

Find k


→ As these 2 lines are \perp to each other,
 (slope of 1st line \times slope of 2nd line) = -1

$$\frac{3k}{22} \times \frac{90}{47} = -1$$

$$270k = -1034$$

$$k = -3.82962962962$$

69.




x intercept of line $7x + 3y = 210$
 is 30 & y intercept of line
 $7x + 3y = 210$ is 70

x intercept of this
 Line is 30
 y intercept of this
 Line is 70

If x intercept of a line is 'm' & y intercept is k then that line passes through $(m, 0)$, $(0, k)$

If x intercept of a line is -20 & y intercept is 35 then that line passes through points:
 $(-20, 0)$ & $(0, 35)$

70. Find Eqⁿ of line having x, y intercepts as a, b respectively.



As x intercept is 'a' & y intercept is 'b', this Line passes through (a,0) & (0,b)

∴ Eqⁿ of the line passing through (a,0) & (0,b) }

$$\frac{b-0}{y-0} = \frac{0-a}{x-a}$$

$$\frac{b}{y} = \frac{-a}{x-a}$$

Intercept form of Eqⁿ of Line

$$b(x-a) = -ay$$

$$bx - ab = -ay$$

$$bx + ay = ab$$

$$\frac{bx}{ab} + \frac{ay}{ab} = \frac{ab}{ab}$$

$$\frac{x}{a} + \frac{y}{b} = 1$$

71. Find Eqⁿ of line passing through points (30, 0), (0, 80)

(30,0) (0,80)

$$\frac{80-0}{y-0} = \frac{0-30}{x-30}$$

$$80(x-30) = -30y$$

$$80x - 2400 = -30y$$

$$80x + 30y = 2400$$

$$8x + 3y = 240$$

OR

$$\text{slope} = \frac{80-0}{0-30}$$

$$= \frac{80}{-30}$$

$$= -8/3$$

Eqⁿ of the line

$$8x + 3y = 240$$

OR

x intercept = 30

y intercept = 80

$$\frac{x}{30} + \frac{y}{80} = 1$$

$$\frac{80x + 30y}{2400} = 1$$

$$80x + 30y = 2400$$

$$8x + 3y = 240$$

72.

Equation of line	x-intercept	y-intercept
$3x + 5y = 90$	30	18
$5x - 2y = 200$	40	-100
$13x + 18y = k$	$\frac{k}{13}$	$\frac{k}{18}$
$20x + 13y = 500$	25	$\frac{500}{13}$
$2x - 11y = -53$	$-\frac{53}{2}$	$\frac{53}{11}$
$21x - y = 200$	$\frac{200}{21}$	-200
$x - y = 10$	10	-10
$2x + y = 58$	29	58
$x = 90$	90	No y-intercept as line is to y-axis
$y = 65$	No x-intercept	65
$kx + my = j$	j/k	j/m
$2kx + 3my = 93$	$93/2k$	$93/3m = 31/m$
$x + 2y = m$	m	$m/2$
$5x + 3y = 1500$	300	500
$x = \frac{90}{7}$	$\frac{90}{7}$	No y-intercept

73. Find eqⁿ of line having x intercept as 3m & y intercept as 38.



Intercept form of Eqⁿ of line is,

$$\frac{x}{a} + \frac{y}{b} = 1 \quad \text{where } a = x \text{ intercept}$$

$$b = y \text{ intercept}$$

$$\left(\frac{x}{3m} + \frac{y}{38} \right) = 1$$

$$\frac{38x + 3my}{114m} = 1 \quad \therefore 38x + 3my = 114m$$

74. Find eqⁿ of line having slope of $\frac{-8}{11}$ and x intercept as 12.



Slope of the line = $-\frac{8}{11}$ \therefore Eqⁿ of Line

Line passes through (12, 0) $8x + 11y = 96$

75. Find slope of the line whose y intercept is 4 times of x intercept.



$a = x$ intercept

$b = y$ intercept = $4a$

$$\frac{x}{a} + \frac{y}{b} = 1$$

Eqⁿ of line having
x intercept as 'a' &
y intercept as 'b'

$$\frac{x}{a} + \frac{y}{4a} = 1$$

$$\frac{4x}{4a} + \frac{y}{4a} = 1$$

$$\frac{4x + y}{4a} = 1$$

$$4x + y = 4a \quad \text{..... slope of this line} = -\frac{4}{1} = -4$$

76. Find slope of the line whose x intercept is $(\frac{4}{5})^{\text{th}}$ of y intercept.



x intercept = $a = \frac{4}{5} b$

y intercept = b

Inequalities & Equations



$$\frac{x}{a} + \frac{y}{b} = 1$$

Eqⁿ of line having
x intercept as 'a' &
y intercept as 'b'

$$\frac{x}{\frac{4b}{5}} + \frac{y}{b} = 1$$

$$\frac{5x}{4b} + \frac{4y}{4b} = 1$$

$$5x + 4y = 4b$$


slope of this line = $-\frac{5}{4}$

Q. Find slope of the line whose x intercept is equal to y intercept.

$$\Rightarrow \frac{x}{a} + \frac{y}{b} = 1 \quad \frac{x}{a} + \frac{y}{a} = 1$$

$\therefore x + y = a = b \quad \therefore$ slope of line = $-\frac{1}{1} = -1$

77. If A = (3,4), O = (0,0). Find l(OA) = ?



$\triangle OAM$ is a Right angle triangle

$$OA^2 = OM^2 + AM^2$$

$$OA^2 = 3^2 + 4^2$$

$$l(OA) = \sqrt{3^2 + 4^2}$$

$$l(OA) = \sqrt{9 + 16}$$

$$l(OA) = \sqrt{25}$$

$$= 5 \text{ units}$$

If O = (0,0), A = (m,n)
then $l(OA) = \sqrt{m^2 + n^2}$

If O \equiv (0,0), K \equiv (12,-5)


Find l(OK)

$$\Rightarrow l(OK) = \sqrt{12^2 + (-5)^2}$$

$$= \sqrt{144 + 25} = 13 \text{ units}$$

A \equiv (0,0), B \equiv (8,12)

$$l(AB) = \sqrt{8^2 + 12^2} = 14.42221 \text{ units}$$



$\triangle PQM$ is a Right angle triangle

$$PQ^2 = PM^2 + QM^2$$

$$PQ^2 = (c-a)^2 + (d-b)^2$$

$$\therefore l(PQ) = \sqrt{(d-b)^2 + (c-a)^2}$$

$$P \equiv (a, b) \quad Q \equiv (c, d)$$

$$\text{then } l(PQ) = \sqrt{(d-b)^2 + (c-a)^2}$$

$P(a, b), Q(c, d)$

$$\text{then } l(PQ) = \sqrt{(d-b)^2 + (c-a)^2}$$

If $A = (x_1, y_1)$ & $B = (x_2, y_2)$ then

$$l(AB) = \sqrt{(y_2 - y_1)^2 + (x_2 - x_1)^2}$$

$$= \sqrt{(y_1 - y_2)^2 + (x_1 - x_2)^2}$$

If $A = (x_1, y_1)$ & $B = (0, 0)$ then

$$l(AB) = \sqrt{x_1^2 + y_1^2}$$

79. $A = (30, 50), B = (80, -90)$. Find $l(AB)$



$$l(AB) = \sqrt{(-90 - 50)^2 + (80 - 30)^2}$$

$$= \sqrt{(-140)^2 + 50^2} = \sqrt{22100} = 148.66069 \text{ units}$$



80. $P = (m,n)$, $Q = (i,j)$ then $l(PQ) =$

$$\begin{aligned} \rightarrow l(PQ) &= \sqrt{(j-n)^2 + (i-m)^2} \\ &= \sqrt{(n-j)^2 + (m-i)^2} \end{aligned}$$

81. If $A = (1.50, 2.875)$, $B = (33, 81.93)$. Find $l(AB)$

$$\begin{aligned} \rightarrow l(AB) &= \sqrt{(81.93 - 2.875)^2 + (33 - 1.50)^2} \\ &= \sqrt{7241.943025} = 85.09961 \text{ units} \end{aligned}$$

82. If $A = (0, 0)$, $B = (-8.75, 33.8175)$. Find $l(AB)$

$$\begin{aligned} \rightarrow l(AB) &= \sqrt{(33.8175 - 0)^2 + (-8.75 - 0)^2} \\ &= \sqrt{1220.18580625} = 34.9312 \text{ units} \end{aligned}$$

83. Points $(8, 13)$, $(16, 19)$, $(-2k, 48)$ are collinear. Find k .

$$\left[\begin{array}{l} \text{slope of the line} \\ \text{passing through} \\ (8, 13) \ \& \ (16, 19) \end{array} \right] = \left[\begin{array}{l} \text{slope of the line} \\ \text{passing through} \\ (-2k, 48), \ (8, 13) \end{array} \right]$$


$$\frac{19-13}{16-8} = \frac{13-48}{8-(-2k)}$$

$$\frac{6}{8} = \frac{-35}{8+2k}$$


$$48 + 12k = -280$$

$$12k = -328$$

$$\therefore k = -27.33333$$



Here points A, B, C are Not collinear



P, Q, R are collinear as a straight line can pass through all of them.

If a straight line can pass through all the points then points are said to be collinear

If a straight line can Not pass through all the points then points are said to be Non collinear

84. Points A, B, C are said to be collinear if

$$\left(\begin{array}{l} \text{Slope of the line passing} \\ \text{through points A, B} \end{array} \right) = \left(\begin{array}{l} \text{Slope of the line passing} \\ \text{through C \& A or B} \end{array} \right)$$

85. Points $(16, \frac{-2k}{5})$, $(8, 11)$, $(19, 85)$ are collinear. Find the value of k.

→ Slope of the line passing through $(16, \frac{-2k}{5})$ & $(8, 11)$ = Slope of the line passing through $(19, 85)$ & $(8, 11)$

$$\frac{11 + \frac{2k}{5}}{8 - 16} = \frac{11 - 85}{8 - 19}$$

$$\frac{55 + 2k}{-8} = \frac{74}{11}$$

$$\therefore \frac{55 + 2k}{5} = \frac{-592}{11}$$

$$55 + 2k = \frac{-2960}{11}$$

$$k = -162.0454545$$

86.

QUADRATIC EQUATION

The standard format of quadratic equation is :

$$ax^2 + bx + c = 0 \quad \text{where } a \neq 0$$

and values of x which can satisfy the quadratic equation are known as 'roots' of quadratic equation.

$x^2 - 10x + 16 = 0$ is a quadratic eqⁿ where $a=1$, $b=-10$, $c=16$

→ Let's put $x=8$ $8^2 - 10(8) + 16 = 0$ Now put $x=2$, $2^2 - 10(2) + 16 = 0$
 $\therefore 8, 2$ are roots of quad. eqⁿ.

Inequalities & Equations



$$x^2 - 5x - 6 = 0 \longrightarrow \text{For } x = 6, x = -1$$

In this quadratic equation $a = 1, b = -5, c = -6$

If we put $x = 6, 6^2 - 5(6) - 6 = 36 - 30 - 6 = 0$

If we put $x = -1, (-1)^2 - 5(-1) - 6 = 1 + 5 - 6 = 0$

this Eqⁿ is satisfied



$\therefore 6, -1$ are roots of quadratic equation $x^2 - 5x - 6 = 0$

87.	Equation	No. of roots
	Linear	1
	Quadratic	2
	Cubic	3

88.

Quadratic Eq ⁿ	a	b	c
$3x^2 + 5x - 8 = 0$	3	5	-8
$19x^2 - 55mx - 2k - 81 = 0$	19	-55m	-2k-81
$15x^2 - 21x - 8px + 39x + 88k - 93 = 18x^2$	-3	$(-21 - 8p + 39)$	$(88k - 93)$
i.e. $-3x^2 + (-21 - 8p + 39)x + 88k - 93 = 0$		$= (18 - 8p)$	
$10x^2 - 2p + 63 = 0$	10	0	-2p+63
$55x^2 - kx^2 + 8px - 33mx + 18j = 63$	$(55 - k)$	$(8p - 33m)$	$(18j - 63)$
i.e. $(55 - k)x^2 + (8p - 33m)x + 18j - 63 = 0$			
$17x^2 - 3x - 93 = 0$	17	-3	-93
$x^2 - 25 = 0$	1	0	-25
$x^2 = 58$	1	0	-58
$(p+q)x^2 - p^2q^2x - 33m = 80$	$(p+q)$	$-p^2q^2$	$(-33m - 80)$



89. Find roots of quadratic equation

$$x^2 - 13x + 36 = 0$$

Formula Method	Short-cut	Super Short-cut
$x^2 - 13x + 36 = 0$	$x^2 - 13x + 36 = 0$	is applicable <u>only when</u>
$a = 1, b = -13, c = 36$	First find the value of	$a = 1$
$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$	$ac = 36$ & $b = -13$	↓
$x = \frac{-(-13) \pm \sqrt{-13^2 - 4(1)(36)}}{2 \times 1}$	Find 2 numbers such	$x^2 - 13x + 36 = 0$
$x = \frac{13 \pm \sqrt{169 - 144}}{2}$	that their sum is 'b'	Find 2 numbers
$x = \frac{13 \pm 5}{2}$	& product is 'ac'	such that their
$x = \frac{13+5}{2}$ OR $x = \frac{13-5}{2}$	$x^2 - 9x - 4x + 36 = 0$	sum is 'b' &
$x = 9$ OR $x = 4$	$x(x-9) - 4(x-9) = 0$	product is 'c'
$\therefore 9, 4$ are the roots	$(x-9)(x-4) = 0$	$(x-9)(x-4) = 0$
	$x-9 = 0$ OR $x-4 = 0$	$x = 9$ / $x = 4$
	$x = 9$ OR $x = 4$	$\therefore 9, 4$ are the roots
	$\therefore 9, 4$ are the roots	of quad. eqn.
	$\therefore 9, 4$ are the roots	
	of quad. eqn.	

90. Find roots of quadratic equation $5x^2 - 13x - 18 = 0$

Formula Method	Short-cut
$a = 5, b = -13, c = -18$	$5x^2 - 13x - 18 = 0$
$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$	$ac = -90, b = -13$
$= \frac{-(-13) \pm \sqrt{(-13)^2 - 4(5)(-18)}}{2 \times 5}$	$5x^2 - 18x + 5x - 18 = 0$
$= \frac{13 \pm \sqrt{529}}{10} = \frac{13+23}{10}$ OR $\frac{13-23}{10}$	$x(5x-18) + 1(5x-18) = 0$
$\therefore x = 18/5$ OR $x = -1$	$(5x-18)(x+1) = 0$
	$\therefore x = 18/5$ OR $x = -1$

$\therefore 18/5, -1$ are the roots of quad. eqn.

91. Find the roots of quadratic equation $2x^2 + 21x + 9 = 0$



$$2x^2 + 21x + 9 = 0$$

This question can not be solved by short-cut.

Let's use formula,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-21 \pm \sqrt{21^2 - 4(2)(9)}}{2 \times 2}$$

$$x = \frac{-21 \pm \sqrt{369}}{4} \quad \therefore x = \frac{-21 + \sqrt{369}}{4} \quad \text{OR} \quad x = \frac{-21 - \sqrt{369}}{4}$$

$$x = -0.4476568 \quad \text{OR} \quad x = \frac{-40.2093}{4}$$

$$x = -10.0523$$

92. Find the roots of $x^2 - 11x - 102 = 0$



$$x = \frac{-(-11) \pm \sqrt{121 - 4(1)(-102)}}{2 \times 1}$$

$$x = \frac{11 \pm 23}{2}$$

$$x = \frac{11+23}{2}, x = \frac{11-23}{2}$$

$$x = 17, x = -6$$

$$x^2 - 11x - 102 = 0$$

$$x^2 - 17x + 6x - 102 = 0$$

$$x(x-17) + 6(x-17) = 0$$

$$(x-17)(x+6) = 0$$

$$x = 17, x = -6$$

$$x^2 - 11x - 102 = 0$$

$$(x-17)(x+6) = 0$$

$$x = 17, x = -6$$

93. Find the roots of $10x^2 - x - 24 = 0$



$$10x^2 - x - 24 = 0$$

$$10x^2 - 16x + 15x - 24 = 0$$

$$2x(5x-8) + 3(5x-8) = 0$$

$$(5x-8)(2x+3) = 0$$

$$x = \frac{8}{5} \quad \text{OR} \quad x = -\frac{3}{2}$$

$$\text{Roots are : } \frac{8}{5}, -\frac{3}{2}$$

$$10x^2 - x - 24 = 0$$

By Formula

$$x = \frac{-(-1) \pm \sqrt{(-1)^2 - 4(10)(-24)}}{2 \times 10}$$

$$x = \frac{1 \pm \sqrt{961}}{20} \quad x = \frac{1+31}{20} \quad \text{OR} \quad \frac{1-31}{20}$$

$$\therefore x = \frac{32}{20} = \frac{8}{5} \quad \text{OR} \quad x = \frac{-30}{20} = -\frac{3}{2}$$

94. Find roots of quadratic equation $80x^2 - 138x + 13 = 0$



$$80x^2 - 138x + 13 = 0$$

$$x = \frac{-(-138) \pm \sqrt{19044 - 4(80)(13)}}{2 \times 80}$$

$$80x^2 - 130x - 8x + 13 = 0$$

$$x = \frac{138 \pm \sqrt{14884}}{160} = \frac{138 \pm 122}{160}$$

$$10x(8x - 13) - 1(8x - 13) = 0$$

$$(8x - 13)(10x - 1) = 0$$

$$x = \frac{138 + 122}{160} \text{ or } x = \frac{138 - 122}{160}$$

$$x = \frac{13}{8} \text{ or } x = \frac{1}{10}$$

$$x = \frac{260}{160} \text{ or } x = \frac{16}{160}$$

Roots are $\frac{13}{8}, \frac{1}{10}$

$$x = \frac{13}{8} \text{ or } x = \frac{1}{10}$$

(i.e. 1.625, 0.10)

95. Find roots of quadratic equation $14x^2 + 29x - 15 = 0$

Also find sum of roots, product of roots.



$$14x^2 + 29x - 15 = 0$$

$$14x^2 + 35x - 6x - 15 = 0$$

$$7x(2x + 5) - 3(2x + 5) = 0$$

$$(2x + 5)(7x - 3) = 0$$

$$\therefore x = -\frac{5}{2} \text{ OR } x = \frac{3}{7}$$

$$\therefore \text{Roots are } -\frac{5}{2} \text{ \& } \frac{3}{7}$$

Sum of roots

$$= \text{1st root} + \text{2nd root}$$

$$= \frac{-5}{2} + \frac{3}{7} = \frac{-35 + 6}{7 \times 2}$$

$$= -\frac{29}{14}$$

Product of roots

$$= \text{1st root} \times \text{2nd root}$$

$$= \frac{-5}{2} \times \frac{3}{7} = -\frac{15}{14}$$

$$14x^2 + 29x - 15 = 0$$

For a Quadratic Eqⁿ

Sum of roots

Product of roots

$$= \frac{-b}{a} = -\frac{29}{14}$$

$$= \frac{c}{a} = \frac{-15}{14}$$

96. Find roots of quadratic equation $10x^2 - 59x - 6 = 0$

Also find sum of roots, product of roots.

$\rightarrow 10x^2 - 59x - 6 = 0$
 $10x^2 - 60x + x - 6 = 0$
 $10x(x-6) + 1(x-6) = 0$
 $(x-6)(10x+1) = 0$
 $x = 6 \text{ OR } x = -\frac{1}{10}$

Roots are : $6, -\frac{1}{10}$

sum of roots = $6 + \frac{-1}{10} = \frac{60-1}{10} = \frac{59}{10}$

product of roots = $6 \times -\frac{1}{10} = \frac{-6}{10} = -\frac{3}{5}$

sum of roots = $-\frac{b}{a}$
 $= -\frac{-59}{10} = \frac{59}{10}$

product of roots = $\frac{c}{a}$
 $= \frac{-6}{10} = -\frac{3}{5}$

97.

Quadratic Equation	Sum of Roots	Product of Roots
$ax^2 + bx + c = 0$	$-\frac{b}{a}$	$\frac{c}{a}$
$8x^2 - 15x - 33 = 0$	$15/8$	$-33/8$
$2x^2 - px + mq + 93 = 0$	$p/2$	$\frac{(mq+93)}{2}$
$x^2 - 40 = 0$	Zero	$\frac{-40}{1} = -40$
$px^2 + qx + r = 0$	$-\frac{q}{p}$	$\frac{r}{p}$
$(3k+3)x^2 - (2p-q)x + 8j + 63 = 0$	$\frac{(2p-q)}{(3k+3)}$	$\frac{(8j+63)}{(3k+3)}$



98. In a Quadratic Equation $ax^2 + bx + c = 0$

1. Sum of roots = 1st root + 2nd root

$$\begin{aligned} \rightarrow &= \left(\frac{-b + \sqrt{b^2 - 4ac}}{2a} \right) + \left(\frac{-b - \sqrt{b^2 - 4ac}}{2a} \right) \\ &= \frac{-b + \sqrt{b^2 - 4ac} - b - \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{-2b}{2a} \\ &= (-b/a) \end{aligned}$$

2. Product of roots = 1st root x 2nd root

$$\begin{aligned} \rightarrow &= \left(\frac{-b + \sqrt{b^2 - 4ac}}{2a} \right) \times \left(\frac{-b - \sqrt{b^2 - 4ac}}{2a} \right) \\ &= \frac{(-b + \sqrt{b^2 - 4ac}) \times (-b - \sqrt{b^2 - 4ac})}{2a \times 2a} \\ &= \frac{(-b)^2 - (\sqrt{b^2 - 4ac})^2}{2a \times 2a} \\ &= \frac{b^2 - (b^2 - 4ac)}{4 \times a \times a} = \frac{\cancel{b^2} - \cancel{b^2} + 4ac}{4a \times a} \\ &= \frac{\cancel{4a} \times c}{\cancel{4a} \times a} = (c/a) \end{aligned}$$

99. If α, β are roots of quadratic equation $5x^2 - 3x - 8 = 0$.

Find the value of $(\alpha + \beta), \alpha\beta, (\alpha + \beta)^2, (\alpha^2 + \beta^2)$

→ 1. $\alpha + \beta = \text{sum of roots} = \frac{3}{5}$

2. $\alpha\beta = \text{product of roots} = \frac{-8}{5}$

3. $(\alpha + \beta)^2 = \left(\frac{3}{5}\right)^2 = \frac{9}{25}$

4. $(\alpha^2 + \beta^2) = (\alpha + \beta)^2 - 2\alpha\beta = \left(\frac{3}{5}\right)^2 - 2\left(\frac{-8}{5}\right) = \frac{9}{25} + \frac{80}{25} = \frac{89}{25}$

$$(\alpha + \beta)^2 = \alpha^2 + \beta^2 + 2\alpha\beta$$

$$(\alpha^2 + \beta^2) = (\alpha + \beta)^2 - 2\alpha\beta$$

$$(\alpha^3 + \beta^3) = (\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)$$

$$(\alpha - \beta)^2 = (\alpha + \beta)^2 - 4\alpha\beta$$

please remember

100. If p, q are roots of quadratic equation $x^2 - 3x + 20 = 0$. Find values of

a. $p + q = -(-3)/1 = 3 = \text{sum of roots}$

b. $pq = \text{product of roots} = 20/1 = 20$

c. $(p - q)^2 = (p + q)^2 - 4pq = 3^2 - 4(20) = -71$

d. $(p^2 + q^2) = (p + q)^2 - 2pq = 3^2 - 2(20) = -31$

e. $p^3 + q^3 = (p + q)^3 - 3pq(p + q) = 3^3 - 3 \times 20(3) = 27 - 180 = -153$

f. $p^2q + q^2p = pq(p + q) = 20 \times 3 = 60$

101. If α, β are roots of quadratic equation $3x^2 - 5x + 2 = 0$.

Find the values of :

$$1. \alpha + \beta = \text{sum of roots} = \left(\frac{5}{3}\right)$$

$$2. \alpha\beta = \text{product of roots} = \left(\frac{2}{3}\right)$$

$$3. (\alpha + \beta)^3 = \left(\frac{5}{3}\right)^3 = \left(\frac{125}{27}\right)$$

$$4. (\alpha^2 + \beta^2) = (\alpha + \beta)^2 - 2\alpha\beta = \frac{25}{9} - 2 \times \frac{2}{3} = \frac{25}{9} - \frac{12}{9} = \left(\frac{13}{9}\right)$$

$$5. (\alpha^3 + \beta^3) = (\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta) \\ = \frac{125}{27} - \left(3 \times \frac{2}{3} \times \frac{5}{3}\right) = \frac{125}{27} - \frac{90}{27} = \left(\frac{35}{27}\right)$$

$$6. (\alpha - \beta)^2 = (\alpha + \beta)^2 - 4\alpha\beta \\ = \frac{25}{9} - 4\left(\frac{2}{3}\right) = \frac{25}{9} - \frac{24}{9} = \left(\frac{1}{9}\right)$$

$$7. \alpha^2\beta + \beta^2\alpha = \alpha\beta(\alpha + \beta) \\ = \frac{2}{3} \times \frac{5}{3} = \left(\frac{10}{9}\right)$$

$$8. \frac{\alpha}{\beta} + \frac{\beta}{\alpha} = \frac{\alpha^2 + \beta^2}{\alpha\beta} = \frac{13/9}{2/3} = \frac{13/9}{6/9} = \left(\frac{13}{6}\right)$$

$$9. \frac{\alpha^2}{\beta} + \frac{\beta^2}{\alpha} = \frac{\alpha^3 + \beta^3}{\alpha\beta} = \frac{35/27}{2/3} = \frac{35/27}{18/27} = \left(\frac{35}{18}\right)$$

102. If α, β are roots of quadratic equation $x^2 + 5x + 13 = 0$.

Find the value of :

1. $\alpha + \beta = -5$

2. $\alpha\beta = 13$

3. $(\alpha^2 + \beta^2) = (\alpha + \beta)^2 - 2\alpha\beta = (-5)^2 - 2(13) = 25 - 26 = -1$

4. $(\alpha + \beta)^2 = (-5)^2 = 25$

5. $(\alpha - \beta)^2 = (\alpha + \beta)^2 - 4\alpha\beta = (-5)^2 - 4(13) = -27$

6. $\alpha^2\beta^2 = (\alpha\beta)^2 = 13^2 = 169$

7. $\alpha^2\beta + \beta^2\alpha = \alpha\beta(\alpha + \beta) = 13 \times -5 = -65$

8. $\frac{\alpha}{\beta} + \frac{\beta}{\alpha} = \left(\frac{\alpha^2 + \beta^2}{\alpha\beta}\right) = \frac{-1}{13} = -\left(\frac{1}{13}\right) = \left(\frac{1}{-13}\right)$

9. $\frac{\alpha^2}{\beta} + \frac{\beta^2}{\alpha} = \frac{\alpha^3 + \beta^3}{\alpha\beta} = \frac{70}{13}$

10. $\alpha^3 + \beta^3 = (\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta) = (-5)^3 - 3(13)(-5)$
 $= -125 + 195$
 $= 70$

103. The standard format of quadratic equation is

$$ax^2 + bx + c = 0 \quad \text{where } a \neq 0$$

$$ax^2 - (-b)x + c = 0$$

dividing by 'a' on both sides

$$\frac{ax^2}{a} - \frac{(-b)}{a}x + \frac{c}{a} = \frac{0}{a}$$

$$x^2 - \frac{(-b)}{a}x + \frac{c}{a} = 0$$

$$x^2 - (\text{sum of roots})x + (\text{product of roots}) = 0$$

$$x^2 - (\text{Sum of roots})x + (\text{Product of roots}) = 0$$

104.

Find roots of quadratic equation

$$10x^2 + 11x + 1 = 0$$



$$10x^2 + 10x + x + 1 = 0$$

$$10x(x+1) + 1(x+1) = 0$$

$$(x+1)(10x+1) = 0$$

$$\therefore x = -1, x = -\frac{1}{10}$$

$\therefore -1, -\frac{1}{10}$ are the roots of quad. eqn.

Find the quadratic equation whose

roots are -1 & $-\frac{1}{10}$



$$x^2 - (\text{sum of roots})x + (\text{product of roots}) = 0$$

$$x^2 - \left(-1 + -\frac{1}{10}\right)x + \left(-1 \times -\frac{1}{10}\right) = 0$$

$$x^2 - \left(-\frac{11}{10}\right)x + \frac{1}{10} = 0$$

$$x^2 + \frac{11}{10}x + \frac{1}{10} = 0$$

$$10x^2 + 11x + 1 = 0$$

105. Find roots of quadratic

equation $16x^2 + 36x - 10 = 0$



$$16x^2 + 36x - 10 = 0$$

$$8x^2 + 18x - 5 = 0$$

$$8x^2 + 20x - 2x - 5 = 0$$

$$4x(2x+5) - 1(2x+5) = 0$$

$$(2x+5)(4x-1) = 0$$

$$x = -\frac{5}{2}, x = \frac{1}{4}$$

Roots are : $-\frac{5}{2}$ & $\frac{1}{4}$

Find the quadratic equation whose

roots are $\frac{1}{4}$ & $-\frac{5}{2}$



$$x^2 - (\text{sum of roots})x + (\text{product of roots}) = 0$$

$$x^2 - \left(\frac{1}{4} + -\frac{5}{2}\right)x + \left(\frac{1}{4} \times -\frac{5}{2}\right) = 0$$

$$x^2 - \frac{-18}{8}x - \frac{5}{8} = 0$$

$$8x^2 + 18x - 5 = 0$$

106. Find roots of quadratic

equation $6x^2 + 19x - 7 = 0$



$$6x^2 + 19x - 7 = 0$$

$$6x^2 + 21x - 2x - 7 = 0$$

$$3x(2x+7) - 1(2x+7) = 0$$

$$(2x+7)(3x-1) = 0$$

$$\therefore x = -\frac{7}{2}, x = \frac{1}{3}$$

Roots are : $\frac{1}{3}$ & $-\frac{7}{2}$

Find the quadratic equation whose

roots are $\frac{1}{3}$ & $-\frac{7}{2}$



$$x^2 - (\text{sum of roots})x + (\text{product of roots}) = 0$$

$$x^2 - \left(\frac{1}{3} - \frac{7}{2}\right)x + \left(\frac{1}{3} \times -\frac{7}{2}\right) = 0$$

$$x^2 - \frac{-19}{6}x - \frac{7}{6} = 0$$

$$6x^2 + 19x - 7 = 0$$

Inequalities & Equations



107. Find roots of quadratic equation $x^2 - 10x + 23 = 0$

→ $x^2 - 10x + 23 = 0$

By Formula,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$= \frac{-(-10) \pm \sqrt{100 - 4(1)(23)}}{2 \times 1}$$

$$= \frac{10 \pm \sqrt{8}}{2} = \frac{10 \pm \sqrt{4 \times 2}}{2}$$

$$= \frac{10 \pm 2\sqrt{2}}{2} = \frac{2(5 \pm \sqrt{2})}{2}$$

$$= (5 \pm \sqrt{2}) \therefore \text{Roots are: } 5 + \sqrt{2} \text{ \& } 5 - \sqrt{2}$$

Find the quadratic equation whose roots are $(5 + \sqrt{2})$ & $(5 - \sqrt{2})$



$$x^2 - (\text{sum of roots})x + (\text{product of roots}) = 0$$

$$x^2 - [5 + \sqrt{2} + 5 - \sqrt{2}]x + [(5 + \sqrt{2})(5 - \sqrt{2})] = 0$$

$$x^2 - 10x + (5^2 - \sqrt{2}^2) = 0$$

$$x^2 - 10x + 23 = 0$$

108.

Roots of Quadratic Eq ⁿ	Quadratic Equation
5, 10	$x^2 - 15x + 50 = 0$
-18, 20	$x^2 - 2x - 360 = 0$
1, -1	$x^2 - 0x - 1 = 0$ i.e. $x^2 - 1 = 0$
15, 18	$x^2 - 33x + 270 = 0$
-16, -20	$x^2 + 36x + 320 = 0$
$-\frac{5}{2}, \frac{9}{2}$	$x^2 - \left(\frac{9}{2} - \frac{5}{2}\right)x + \left(\frac{9}{2} \times -\frac{5}{2}\right) = 0$ $x^2 - 2x - \frac{45}{4} = 0, 4x^2 - 8x - 45 = 0$
$\frac{9}{7}, \frac{8}{13}$	$x^2 - \left(\frac{9}{7} + \frac{8}{13}\right)x + \left(\frac{9}{7} \times \frac{8}{13}\right) = 0$ $x^2 - \left(\frac{173}{91}\right)x + \frac{72}{91} = 0, 91x^2 - 173x + 72 = 0$
16, 0	$x^2 - 16x = 0$
$(8 + \sqrt{3}), (8 - \sqrt{3})$	$x^2 - (16)x + (61) = 0$
$(1 + \sqrt{30}), (1 - \sqrt{30})$	$x^2 - (2)x + (-29) = 0$ $x^2 - 2x - 29 = 0$

109. Find roots of quadratic equation $4x^2 + 12x + 9 = 0$



$$4x^2 + 12x + 9 = 0$$

$$4x^2 + 6x + 6x + 9 = 0$$

$$2x(2x+3) + 3(2x+3) = 0$$

$$(2x+3)(2x+3) = 0$$

$$\therefore 2x+3 = 0 \quad \text{OR} \quad 2x+3 = 0$$

$$x = -\frac{3}{2} \quad \text{OR} \quad x = -\frac{3}{2}$$

\therefore Roots are $-\frac{3}{2}$ & $-\frac{3}{2}$

$$x = \frac{-12 \pm \sqrt{12^2 - 4(4)(9)}}{2 \times 4}$$

$$x = \frac{-12 \pm 0}{8}$$

$$x = \frac{-12+0}{8} \quad \text{OR} \quad \frac{-12-0}{8}$$

$$x = -\frac{3}{2} \quad \text{OR} \quad x = -\frac{3}{2}$$

$$b^2 - 4ac = 12^2 - 4(4)(9) = 144 - 144 = 0$$

When $b^2 - 4ac = 0$ then roots of quadratic equation are equal.

110. Find roots of quadratic equation $5x^2 + 15x + 91 = 0$



Let's solve by formula

$$x = \frac{-15 \pm \sqrt{15^2 - 4(5)(91)}}{2 \times 5}$$

$$x = \frac{-15 \pm \sqrt{225 - 1820}}{10}$$

$$x = \frac{-15 \pm \sqrt{-1595}}{10}$$

Roots are imaginary/unreal/complex numbers

When $(b^2 - 4ac) < 0$ then roots of quadratic equation are unreal/complex/Imaginary numbers

111. Find roots of quadratic equation $x^2 - 14x + 46 = 0$



$$x^2 - 14x + 46 = 0$$

Let's use formula

$$x = \frac{-(-14) \pm \sqrt{(-14)^2 - 4(1)(46)}}{2 \times 1} = \frac{14 \pm \sqrt{196 - 184}}{2}$$

$$x = \frac{14 \pm \sqrt{12}}{2} = \frac{14 \pm \sqrt{4 \times 3}}{2}$$

$$x = \frac{14 \pm 2\sqrt{3}}{2} = \frac{2(7 \pm \sqrt{3})}{2} = (7 \pm \sqrt{3})$$

Roots are : $(7 + \sqrt{3})$ & $(7 - \sqrt{3})$

we can clearly see that : Roots are Irrational

when $(b^2 - 4ac) > 0$ & Not a perfect square
then Roots of quadratic eqⁿ are Irrational

Find roots of : $3x^2 - 8x - 11 = 0$

$$\Rightarrow x = \left[\frac{-(-8) \pm \sqrt{(-8)^2 - 4(3)(-11)}}{2 \times 3} \right] = \frac{8 \pm \sqrt{196}}{6}$$

$$= \frac{8+14}{6} \text{ OR } \frac{8-14}{6} = \frac{11}{3} \text{ OR } -1$$

when $(b^2 - 4ac) > 0$ & a perfect square
then Roots of quadratic eqⁿ are Rational

112.

When	Nature of roots
$b^2 - 4ac = 0$	Real, Rational, Equal
$b^2 - 4ac < 0$	Unreal / imaginary / complex
$b^2 - 4ac > 0$ and not a perfect square	Real, Irrational, unequal
$b^2 - 4ac > 0$ and a perfect square	Real, Rational, unequal

value of $b^2 - 4ac$	Nature of roots
38	Real, Irrational, unequal
36	Real, Rational, unequal
81	Real, Rational, unequal
90	Real, Irrational, unequal
-144	Complex / imaginary / unreal
0	Real, Rational, Equal
207936	Real, Rational, unequal
810	Real, Irrational, unequal
-90	Complex / imaginary / unreal
-35	Complex / imaginary / unreal
-0	Real, Rational, Equal
905	Real, Irrational, unequal
2025	Real, Rational, unequal
86	Real, Irrational, unequal
100	Real, Rational, unequal

Quadratic Equation	$b^2 - 4ac$	Nature of roots
$3x^2 - 12x + 1 = 0$	$(-12)^2 - 4(3)(1)$ $= 144 - 12 = 132$	Real, Irrational, Distinct
$5x^2 - 12x = 0$	$(-12)^2 - 4(5)(0)$ $= 144 - 0 = 144$	Real, Rational, unequal
$4x^2 + 12x + 9 = 0$	$12^2 - 4(4)(9)$ $= 144 - 144 = 0$	Real, Rational, Equal
$x^2 - 10x + 23 = 0$	$(-10)^2 - 4(1)(23)$ $= 100 - 92 = 8$	Real, Irrational, Distinct
$10x^2 - x - 9 = 0$	$(-1)^2 - 4(10)(-9)$ $= 1 + 360 = 361$	Real, Rational, unequal
$8x^2 + 11 = 0$	$0^2 - 4(8)(11)$ $= -352$	complex / imaginary / unequal

113. Find the quadratic equation whose roots are $\frac{3}{2}, \frac{-8}{11}$

→ $x^2 - (\text{sum of roots})x + (\text{product of roots}) = 0$

$$x^2 - \left(\frac{3}{2} + \frac{-8}{11}\right)x + \left(\frac{3}{2} \times \frac{-8}{11}\right) = 0$$

$$x^2 - \left(\frac{17}{22}\right)x + \frac{-24}{22} = 0$$

$$22x^2 - 17x - 24 = 0$$

114. Find the quadratic equation whose roots are $(2 + \sqrt{23})$ & $(2 - \sqrt{23})$

→ $x^2 - (\text{sum of roots})x + (\text{product of roots}) = 0$

$$x^2 - (2 + \sqrt{23} + 2 - \sqrt{23})x + [(2 + \sqrt{23})(2 - \sqrt{23})] = 0$$

$$x^2 - 4x - 19 = 0$$

115. Find the quadratic equation whose one root is $(15 + \sqrt{41})$

→ If one root of quad. eqⁿ is $(15 + \sqrt{41})$ then other root must be $(15 - \sqrt{41})$

∴ Roots are : $(15 + \sqrt{41})$ & $(15 - \sqrt{41})$

∴ Quadratic Eqⁿ is :

$$x^2 - (\text{sum of roots})x + (\text{product of roots}) = 0$$

$$x^2 - 30x + (225 - 41) = 0$$

$$x^2 - 30x + 184 = 0$$

116.

Quadratic Equation	$b^2 - 4ac$	Nature of Roots
$3x^2 - 5x - 8 = 0$	$(-5)^2 - 4(3)(-8)$ $= 121$	Real, Rational, unequal
$8x^2 - 13x + 200 = 0$	$(-13)^2 - 4(8)(200)$ $= 169 - 6400$ $= -6231$	unreal / complex / Imaginary
$5x^2 + 11x - 3 = 0$	$11^2 - 4(5)(-3)$ $= 121 + 60 = 181$	Real, Irrational, unequal
$4x^2 + 12x + 9 = 0$	$12^2 - 4(4)(9)$ $= 0$	Real, Rational, Equal
$x^2 - 13x + 36 = 0$	$(-13)^2 - 4(1)(36)$ $= 25$	Real, Rational, unequal
$5x^2 + 12x + 7 = 0$	$12^2 - 4(5)(7)$ $= 4$	Real, Rational, unequal
$4x^2 - 1 = 0$	$0^2 - 4(4)(-1)$ $= 16$	Real, Rational, unequal
$3x^2 + 22x = 0$	$22^2 - 4(3)(0)$ $= 484$	Real, Rational, unequal
$8x^2 - 2x + 33 = 0$	$(-2)^2 - 4(8)(33)$ $= 4 - 1056 = -1052$	unreal / complex / Imaginary

117.	Value of $b^2 - 4ac$	Nature of Roots
	38	Real, Irrational, unequal
	41	Real, Irrational, unequal
	49	Real, Rational, unequal
	-60	unreal/complex/Imaginary
	0	Real, Rational, Equal
	88	Real, Irrational, unequal
	14641	Real, Rational, unequal
	19288	Real, Irrational, unequal
	3364	Real, Rational, unequal
	-0	Real, Rational, Equal
	380	Real, Irrational, unequal
	-100	unreal/complex/Imaginary

118. Roots of quadratic equation $5x^2 - 33x + 8k + 5 = 0$ are equal. Find k.

→ AS Roots of quadratic eqn are equal

$$\therefore b^2 - 4ac = 0$$

$$(-33)^2 - 4(5)(8k+5) = 0 \quad \therefore k = \left(\frac{989}{160}\right)$$

$$1089 - 20(8k+5) = 0$$

$$1089 - 160k - 100 = 0$$

$$989 = 160k$$

$$= 6.18125$$

119. Roots of quadratic equation $5kx^2 - 3x^2 + 18x - 21 = 0$ are equal.

Find k.

$$(5k-3)x^2 + 18x - 21 = 0$$

→ AS Roots of quadratic eqn are equal $\therefore 324 + 420k - 252 = 0$

$$\therefore b^2 - 4ac = 0$$

$$18^2 - 4(5k-3)(-21) = 0$$

$$324 + 84(5k-3)$$

$$420k = -72$$

$$k = \frac{-72}{420} = \frac{-18}{105}$$

$$k = -6/35$$

120. Roots of quadratic equation $5mx^2 + 33x - 28 = 0$ are equal. Find m .

→ AS Roots of quadratic eqn are equal

$$\therefore b^2 - 4ac = 0$$

$$33^2 - 4(5m)(-28) = 0$$

$$1089 + 560m = 0$$

$$m = \frac{-1089}{560}$$

121. Roots of quadratic equation $5kx^2 - 33x + 8k - 19 = 0$ are reciprocals of each other. Find the value of k .

→ AS Roots of quadratic eqn are reciprocals of each other,

$$\text{1st root} \times \text{2nd root} = 1$$

$$\text{product of roots} = 1$$

$$\frac{c}{a} = 1$$

$$c = a$$

$$a = c$$

$$5k = 8k - 19$$

$$19 = 3k$$

$$\therefore k = \frac{19}{3}$$

122. Roots of quadratic equation $5x^2 - 8kx + 33x - 8p - 19 = 0$ are equal but opposite in sign. Find the value of k .

→ AS Roots are equal but opposite in sign for

$$\text{quad. eqn} : 5x^2 + (-8k + 33)x - 8p - 19 = 0$$

Inequalities & Equations



$$\text{1st root} + \text{2nd root} = 0$$

$$\text{sum of roots} = 0$$

$$\frac{-b}{a} = 0$$

$$-b = 0$$

$$b = 0$$

$$(-8k + 33) = 0$$

$$33 = 8k$$

$$\therefore k = \frac{33}{8}$$

123. If Roots of quadratic equation are

then

Equal

$$b^2 - 4ac = 0$$

Reciprocal of each other

$$a = c$$

Equal but opposite in sign

$$b = 0 = \text{Zero}$$

124. Roots of quadratic equation $5x^2 + kx^2 - 19x - 33k - 93 = 0$ are reciprocal of each other. Find k.

$$\rightarrow (5+k)x^2 - 19x - 33k - 93 = 0$$

As Roots are Reciprocals of each other,

$$a = c$$

$$5+k = -33k - 93$$

$$34k = -98$$

$$k = -\frac{98}{34} = -\frac{49}{17}$$

125. Roots of quadratic equation $5x^2 - 8px + 81x = 93x - 63k + 88$ are equal but opposite in sign. Find p.

$$\rightarrow 5x^2 + (-8p + 81 - 93)x + 63k - 88 = 0$$

As Roots are Equal but opposite in sign,

$$b = 0$$

$$-8p + 81 - 93 = 0$$

$$-12 = 8p$$

$$\therefore p = -\frac{12}{8} = -\frac{3}{2}$$

126. If p, q are roots of quadratic equation $x^2 - 11x - 28 = 0$. Find values.

1. $p + q = 11$ = sum of roots

2. $pq = -28$ = product of roots

3. $p^3 + q^3 = (p+q)^3 - 3pq(p+q) = 11^3 - 3(-28)(11) = 1331 + 924 = 2255$

4. $p^2 + q^2 = (p+q)^2 - 2pq = 11^2 - 2(-28) = 177$

5. $(p - q)^2 = (p+q)^2 - 4pq = 11^2 - 4(-28) = 233$

6. $\frac{p}{q} + \frac{q}{p} = \frac{p^2 + q^2}{pq} = \frac{177}{-28} = -\frac{177}{28}$

7. $\frac{p^2}{q} + \frac{q^2}{p} = \frac{p^3 + q^3}{pq} = \frac{2255}{-28} = -\frac{2255}{28}$

8. $p^2q + q^2p = pq(p+q) = -28 \times 11 = -308$

9. $(p - q) = \sqrt{(p - q)^2} = \sqrt{233}$

10. $p^2q^2 = (pq)^2 = (-28)^2 = 784$

127. If α, β are roots of quadratic equation $5x^2 - 2x + 3 = 0$.

Find quadratic equation whose roots are $(\alpha^2 + \beta^2), (\alpha - \beta)^2$



$$\alpha + \beta = 2/5$$

$$\alpha\beta = 3/5$$

$$\begin{aligned} (\alpha^2 + \beta^2) &= (\alpha + \beta)^2 - 2\alpha\beta = \frac{4}{25} - 2 \times \frac{3}{5} = \frac{4}{25} - \frac{30}{25} \\ &= -\frac{26}{25} = -1.04 \end{aligned}$$

$$\begin{aligned} (\alpha - \beta)^2 &= (\alpha + \beta)^2 - 4\alpha\beta = \frac{4}{25} - 4 \times \frac{3}{5} = \frac{4}{25} - \frac{60}{25} \\ &= -\frac{56}{25} = -2.24 \end{aligned}$$

→ question is:

Find quad. eqn whose roots are -1.04 & -2.24

∴ Answer is: $x^2 - (-1.04 + -2.24)x + (-1.04 \times -2.24) = 0$

$$x^2 + 3.28x + 2.3296 = 0$$

$$10000x^2 + 32800x + 23296 = 0$$

$$625x^2 + 2050x + 1456 = 0$$

128. If α, β are roots of quadratic equation $x^2 - 12x + 17 = 0$.

Find quadratic equation whose roots are $(\alpha^3 + \beta^3), (\alpha^2 + \beta^2)$

→ $\alpha + \beta = 12, \alpha\beta = 17$

$$\alpha^3 + \beta^3 = 1728 - (3 \times 17 \times 12) = 1116$$

$$\alpha^2 + \beta^2 = 144 - 2 \times 17 = 110$$

Find quad. eqn whose roots are 1116 & 110

Answer: $x^2 - (1116 + 110)x + (1116 \times 110) = 0$

$$x^2 - 1226x + 122760 = 0$$

129. If α, β are roots of quadratic equation $5x^2 - 2x - 11 = 0$.

Find quadratic equation whose roots are $(\alpha + \beta), (\alpha\beta)$

→ $\alpha + \beta = \frac{2}{5}, \alpha\beta = -\frac{11}{5}$

Question is: Find quad. eqⁿ whose roots are $\frac{2}{5}$ & $-\frac{11}{5}$

$$x^2 - \left(\frac{2}{5} - \frac{11}{5}\right)x + \left(\frac{2}{5} \times -\frac{11}{5}\right) = 0$$

$$x^2 - \frac{-9}{5}x - \frac{22}{25} = 0$$

$$25x^2 + 45x - 22 = 0$$

130. If $a+b = 12, ab = 60$. Find $\left(\frac{1}{a} + \frac{1}{b}\right) = ?$

→ $\frac{1}{a} + \frac{1}{b} = \frac{a+b}{ab} = \frac{12}{60} = \frac{1}{5}$

131. Find quadratic equation whose one root is $(11 + \sqrt{13})$

→ Roots are: $(11 + \sqrt{13}), (11 - \sqrt{13})$

Quad. Eqⁿ is: $x^2 - [11 + \sqrt{13} + 11 - \sqrt{13}]x + [(11 + \sqrt{13})(11 - \sqrt{13})] = 0$

$$x^2 - 22x + (121 - 13) = 0$$

$$x^2 - 22x + 108 = 0$$

132. Find quadratic equation whose one root is $(7 + \sqrt{230})$

→ Roots are: $(7 + \sqrt{230})$ & $(7 - \sqrt{230})$

Quad. Eqⁿ is: $x^2 - 14x + (7^2 - \sqrt{230}^2) = 0$

$$x^2 - 14x - 181 = 0$$

133. Standard format of a quadratic equation is :

$$ax^2 + bx + c = 0$$

$$x^2 - (\text{sum of roots})x + (\text{product of roots}) = 0$$

Where $a \neq 0$

134.

Roots of Quadratic Eq ⁿ	Factors of Quadratic Eq ⁿ	Quadratic Eq ⁿ
3, -13	$(x-3), (x+13)$	$x^2 + 10x - 39 = 0$
$-\frac{3}{2}, \frac{1}{8}$	$(2x+3)(8x-1)$	$16x^2 - 2x + 24x - 3 = 0$ $16x^2 + 22x - 3 = 0$
$\frac{1}{2}, \frac{1}{2}$	$4x^2 - 2x - 2x + 1 = 0$ $2x(2x-1) - 1(2x-1) = 0$ $(2x-1) \& (2x-1)$	$4x^2 - 4x + 1 = 0$
$\frac{2}{5}, \frac{9}{8}$	$(5x-2), (8x-9)$	$40x^2 - 45x - 18x + 18 = 0$ $40x^2 - 63x + 18 = 0$
$-\frac{3}{5}, \frac{7}{11}$	$(5x+3), (11x-7)$	$55x^2 - 35x + 33x - 21 = 0$ $55x^2 - 2x - 21 = 0$
$-\frac{4}{3}, -\frac{1}{2}$	$(3x+4), (2x+1)$	$6x^2 + 3x + 8x + 4 = 0$ $6x^2 + 11x + 4 = 0$
$\frac{7}{5}, -\frac{11}{8}$	$(5x-7), (8x+11)$	$40x^2 + 55x - 56x - 77 = 0$ $40x^2 - x - 77 = 0$
0, 8	$x, (x-8)$	$x^2 - 8x = 0$
1, -1	$(x-1), (x+1)$	$x^2 - 1 = 0$
$\frac{5}{3}, \frac{3}{5}$	$(3x-5), (5x-3)$	$15x^2 - 9x - 25x + 15 = 0$ $15x^2 - 34x + 15 = 0$

135. Standard format of a cubic eqⁿ $ax^3 + bx^2 + cx + d = 0$, where $a \neq 0$
 3 values of x can satisfy cubic equation

\therefore Cubic equation has 3 roots

136. Find cubic eqⁿ whose roots are 8, 3, -2.

→ cubic eqⁿ is,

$$(x-8)(x-3)(x+2) = 0$$

$$(x^2 - 11x + 24)(x+2) = 0$$

$$x^3 + 2x^2 - 11x^2 - 22x + 24x + 48 = 0$$

$$\therefore x^3 - 9x^2 + 2x + 48 = 0$$

OR

$$x^3 - (\text{sum of roots})x^2 +$$

$$\left[(1^{\text{st}} \times 2^{\text{nd}}) + (2^{\text{nd}} \times 3^{\text{rd}}) + (1^{\text{st}} \times 3^{\text{rd}}) \right] x - (\text{product of roots}) = 0$$

$$x^3 - (8+3-2)x^2 + (24-16-6)x - (8 \times 3 \times -2) = 0$$

$$x^3 - 9x^2 + 2x + 48 = 0$$

137. Find cubic eqⁿ whose roots are p, q, r

→ AS Roots of cubic eqⁿ are p, q, r

\therefore Factors must be $(x-p), (x-q), (x-r)$

\therefore The cubic Eqⁿ is :

$$(x-p)(x-q)(x-r) = 0$$

$$(x^2 - xq - px + pq)(x-r) = 0$$

$$x^3 - x^2r - x^2q + xqr - px^2 + pxr + pqx - pqr = 0$$

$$x^3 - x^2r - x^2q - x^2p + pqx + qr x + pr x - pqr = 0$$

$$x^3 - (p+q+r)x^2 + (pq+qr+pr)x - (pqr) = 0$$

$$x^3 - (\text{sum of roots})x^2 + \left[(1^{\text{st}} \times 2^{\text{nd}}) + (2^{\text{nd}} \times 3^{\text{rd}}) + (1^{\text{st}} \times 3^{\text{rd}}) \right] x - (\text{product of roots}) = 0$$

compare this with $ax^3 + bx^2 + cx + d = 0$

\therefore Sum of roots = $-b/a$ & product of roots = $-d/a$

138. 1. Find cubic eqⁿ whose roots are 3, -11, -15.

$$\rightarrow x^3 - \left(\text{sum of roots} \right) x^2 + \left[(1^{\text{st}} \times 2^{\text{nd}}) + (2^{\text{nd}} \times 3^{\text{rd}}) + (1^{\text{st}} \times 3^{\text{rd}}) \right] x - \left(\text{product of roots} \right) = 0$$

$$x^3 - (3 - 11 - 15) x^2 + [-33 + 165 + -45] x - (495) = 0$$

$$x^3 + 23x^2 + 87x - 495 = 0$$

2. Find cubic eqⁿ whose roots are m, n, v

$$x^3 - (m+n+v)x^2 + (mn+nv+mv)x - mnv = 0$$

139.

	Cubic Equation	Quadratic Equation
Standard Format	$ax^3 + bx^2 + cx + d = 0$ where $a \neq 0$	$ax^2 + bx + c = 0$ where $a \neq 0$
Sum of roots	$-b/a$	$-b/a$
Product of roots	$-d/a$	c/a

Find sum of roots & product of roots for

$$8x^3 - 3x^2 - 11kx^2 + 3px^3 - 22x + 18kx - 13mx - 2k + 18 = 0$$

\Rightarrow cubic eqⁿ is :

$$(8+3p)x^3 + (-3-11k)x^2 + (-22+18k-13m)x + (-2k+18) = 0$$

$$\therefore \text{sum of roots} = -b/a = \frac{3+11k}{8+3p}$$

$$\text{product of roots} = -d/a = \frac{2k-18}{8+3p}$$

140. Find cubic eqⁿ whose roots are $(\frac{5}{2}, \frac{9}{2}, -\frac{11}{2})$

$$\rightarrow x^3 - \left(\frac{5}{2} + \frac{9}{2} - \frac{11}{2}\right)x^2 + \left[\frac{45}{4} + \frac{-99}{4} + \frac{-55}{4}\right]x - \left(\frac{5}{2} \times \frac{9}{2} \times \frac{-11}{2}\right) = 0$$

$$x^3 - \left(\frac{3}{2}\right)x^2 + \left(\frac{-109}{4}\right)x + \left(\frac{495}{8}\right) = 0$$

$$8x^3 - 12x^2 - 218x + 495 = 0$$

OR

$$(2x-5)(2x-9)(2x+11) = 0$$

$$(2x-5)(4x^2+4x-99) = 0$$

$$8x^3 + 8x^2 - 198x - 20x^2 - 20x + 495 = 0$$

$$8x^3 - 12x^2 - 218x + 495 = 0$$

141. Find quadratic eqⁿ whose roots are $(\frac{5}{2}, 0)$

$$\rightarrow x^2 - (\text{sum of roots})x + (\text{product of roots}) = 0$$

$$x^2 - \frac{5}{2}x + 0 = 0$$

$$2x^2 - 5x = 0$$

OR Factors: $(2x-5)$ & x

$$x(2x-5) = 0 \quad \text{i.e.} \quad 2x^2 - 5x = 0$$

142. Find quadratic eqⁿ whose roots are $(10, -10)$

$$\rightarrow x^2 - (\text{sum of roots})x + (\text{product of roots}) = 0$$

$$x^2 - 0x + (-100) = 0 \quad \therefore x^2 - 100 = 0$$

OR

$$(x-10)(x+10) \text{ are factors} \quad \therefore (x-10)(x+10) = 0$$

$$x^2 - 100 = 0$$

143. Find quadratic eqⁿ whose roots are $(\frac{8}{9}, \frac{9}{8})$

$$\rightarrow x^2 - (\text{sum of roots})x + (\text{product of roots}) = 0$$

$$x^2 - \left(\frac{8}{9} + \frac{9}{8}\right)x + \left(\frac{8}{9} \times \frac{9}{8}\right) = 0$$

$$x^2 - \frac{145}{72}x + 1 = 0 \quad \therefore 72x^2 - 145x + 72 = 0$$

144. The point $(-3p, 28)$ lie on the line $7x + 12y = 820$. Find p .

→ If we put $x = -3p$ & $y = 28$, $7x + 12y = 820$ must be satisfied

$$7(-3p) + 12(28) = 820$$

$$-21p = 484$$

$$p = \left(\frac{-484}{21} \right) = -23.047619$$

145. The point of intersection of lines $7x + 3y = 90$ & $8x + 7y = 210$ lie in _____ Quadrant.

a. 1st

b. 2nd

c. 4th

~~d. None of these~~

→

$$56x + 24y = 720$$

$$56x + 49y = 1470$$

$$-25y = -750$$

$$y = 30$$

$$7x + 3(30) = 90$$

$$7x = 0 \quad \therefore x = 0$$

$(0, 30)$ is the point of intersection.

It is on y -axis

146. The lines $2x + 3y = 90$ & $4x + 6y = 180$ have _____

a. No solution

b. Unique Solution

~~c. Infinite No. of solutions~~

d. None of these

$$2x + 3y = 90 \quad \text{--- ①}$$

$$4x + 6y = 180 \quad \text{--- ②}$$

$$4x + 6y = 180 \quad \left. \begin{array}{l} \text{--- ②} \\ \text{--- ③} \end{array} \right\} \text{These 2 lines coincide}$$

i.e. every point is point of intersection

147. Slope of the line $8x = \frac{81}{11}$ is

a. Zero

b. $\frac{81}{88}$

c. $-\frac{81}{88}$

~~d. Not defined~~

$$8x = \frac{81}{11}$$

$$\therefore 88x + 0y = 81$$

$$88x = 81$$

$$\text{slope} = -a/b = -88/0 = \text{Not defined}$$

$$8x = \frac{81}{11} \text{ is}$$

OR a // line to y -axis

\therefore It's slope is

Not defined

148. Find equation of line having slope of $\left(\frac{8}{11}\right)$ passing through

$\left(\frac{3}{5}, \frac{8}{5}\right)$

→ Slope = $\frac{8}{11}$

∴ Eqⁿ of line $8x - 11y = 8\left(\frac{3}{5}\right) - 11\left(\frac{8}{5}\right)$

$$8x - 11y = -\frac{64}{5}$$

$$40x - 55y = -64$$

$$40x - 55y + 64 = 0$$

149. Find equation of line having x, y intercept as $\frac{8}{3}, \frac{11}{9}$ respectively.

→ Intercept form is $\frac{x}{a} + \frac{y}{b} = 1$

$$\left(\frac{x}{\frac{8}{3}} + \frac{y}{\frac{11}{9}}\right) = 1$$

a = x intercept

b = y intercept

$$\frac{3x}{8} + \frac{9y}{11} = 1$$

$$\frac{33x + 72y}{88} = 1 \quad \therefore 33x + 72y = 88$$

150. Find nature of roots of $3x^2 - 14x - 31 = 0$

→ $b^2 - 4ac = (-14)^2 - 4(3)(-31)$

$$= 196 + 372 = 568 > 0 \text{ but not a}$$

perfect square

∴ Roots are : Real, Irrational, unequal

151. If α, β are roots of quadratic equation $x^2 - 5x + 9 = 0$ then
Find quadratic equation whose roots are $(2\alpha + 3\beta)$ & $(3\alpha + 2\beta)$

→ $\alpha + \beta = 5, \alpha\beta = 9$

quad. eqn whose roots are $(2\alpha + 3\beta)$ & $(3\alpha + 2\beta)$ is

$$x^2 - (\text{sum of roots})x + (\text{product of roots}) = 0$$

$$x^2 - [2\alpha + 3\beta + 3\alpha + 2\beta]x + [(2\alpha + 3\beta)(3\alpha + 2\beta)] = 0$$

$$x^2 - (5\alpha + 5\beta)x + [6\alpha^2 + 4\alpha\beta + 9\alpha\beta + 6\beta^2] = 0$$

$$x^2 - 5(\alpha + \beta)x + [13\alpha\beta + 6(\alpha^2 + \beta^2)] = 0$$

$$x^2 - (5 \times 5)x + [13 \times 9 + 6(5^2 - 2 \times 9)] = 0$$

$$x^2 - 25x + (117 + 6 \times 7) = 0$$

$$x^2 - 25x + (117 + 42) = 0$$

$$x^2 - 25x + 159 = 0$$

152. One root of quadratic equation $3kx^2 + 18px - 19p + 21 = 0$ is 'zero'. Find value of 'p'.

→ one root is zero, means for $x = 0$,
Eqn is satisfied

$$\therefore 3k(0)^2 + 18p(0) - 19p + 21 = 0$$

$$0 + 0 + 21 = 19p$$

$$\therefore p = \frac{21}{19}$$

153. If α, β are roots of quadratic equation $5x^2 - 11x + 29 = 0$,
Find quadratic equation whose roots are $(\alpha + 1)$ & $(\beta + 1)$

→ $\alpha + \beta = \frac{11}{5}, \alpha\beta = \frac{29}{5}$

Quad. Eqⁿ whose roots are $(\alpha+1)$ & $(\beta+1)$ is

$$x^2 - [\alpha + 1 + \beta + 1]x + [(\alpha + 1)(\beta + 1)] = 0$$

$$x^2 - (\alpha + \beta + 2)x + (\alpha\beta + \alpha + \beta + 1) = 0$$

$$x^2 - \left(\frac{11}{5} + 2\right)x + \left(\frac{29}{5} + \frac{11}{5} + \frac{5}{5}\right) = 0$$

$$x^2 - \left(\frac{21}{5}\right)x + \left(\frac{45}{5}\right) = 0$$

$$5x^2 - 21x + 45 = 0$$

154. The points $(16, (-2k/9)), (18, 0), (19, -23)$ are collinear. Find 'k'

→ $\left[\begin{array}{l} \text{Slope of the line} \\ \text{passing through points} \\ (16, -\frac{2k}{9}) \text{ \& } (18, 0) \end{array} \right] = \left[\begin{array}{l} \text{Slope of the line} \\ \text{passing through points} \\ (18, 0) \text{ \& } (19, -23) \end{array} \right]$

$$\frac{0 + \frac{2k}{9}}{18 - 16} = \frac{-23 - 0}{19 - 18}$$

$$\frac{\frac{2k}{9}}{2} = \frac{-23}{1}$$

$$\frac{2k}{9} = -46$$

$$2k = -414$$

$$k = -207$$

155. $\frac{x+24}{5} = 4 + \frac{x}{4}$ Find x.



$$\frac{x+24}{5} = 4 + \frac{x}{4}$$

$$\frac{x+24}{5} = \frac{16+x}{4}$$

$$4x + 96 = 80 + 5x$$

$$16 = x$$

$$\therefore x = 16$$

156. $x + 5y = 36, \frac{x+y}{x-y} = \frac{5}{3}$ then $(x, y) = ?$



$$x = 36 - 5y$$

$$\frac{36 - 5y + y}{36 - 5y - y} = \frac{5}{3}$$

$$\therefore x = 36 - 5(4)$$

$$= 16$$

$$\frac{36 - 4y}{36 - 6y} = \frac{5}{3}$$

$$108 - 12y = 180 - 30y$$


$$18y = 72$$

$$y = 4$$

$$\therefore (x, y) \equiv (16, 4)$$

157. Diagonal of a rectangle is 5 cms and one of the side is 4 cms then

Find Area of Rectangle.




$$5^2 = 4^2 + x^2$$

$$x^2 = 9$$

$$x = 3\text{cms}$$

3cms



4cms

$$\text{Area} = \text{Length} \times \text{Breadth}$$

$$= 4\text{cms} \times 3\text{cms}$$


$$= 12 \text{ sq. cms}$$

158. If one root of quadratic equation exceeds the other by 4 in $x^2 - 8x + m = 0$. Find m.



$$x^2 - (\text{sum of roots})x + \text{product of roots} = 0$$

$$\text{sum of roots} = 8, \quad \text{product of roots} = m$$



\therefore Roots are : 2, 6

$$\therefore \text{product of roots} = 2 \times 6 = 12 = m$$

$\therefore m = 12$

$$x^2 - (2+6)x + (2 \times 6) = 0$$

159. $x + y = 50, \frac{1}{x} + \frac{1}{y} = \frac{1}{8}$ then $(x,y) = ?$



$$x + y = 50 \quad \frac{1}{x} + \frac{1}{y} = \frac{1}{8}$$

$$\frac{x+y}{xy} = \frac{1}{8}$$

$$xy = 8(x+y)$$

$$xy = 8 \times 50 = 400$$

$x + y = 50$



$x \times y = 400$

$\therefore x, y = (10, 40) \text{ OR } (40, 10)$




160. Find feasible area for $x + y \leq 50$

$x + y = 50$	} Linear Equation Or Linear Equality	$x + y \leq 50$	} Linear Inequations Or Linear Inequality
$2x + 3y = 90$		$2x + 3y \geq 90$	
$3x - 5y = 60$		$5x - 18y < 35$	
$x = 35$		$x \leq 48$	
		$y \geq 90$	

$x + y \leq 50$ is a linear inequality.

Let's draw the line $x + y = 50$ on graph paper.



Graphical representation of a linear equation is :
a Line


Graphical representation of a linear inequality is :
AN AREA

Feasible Area or Solution Space for $x + y \leq 50$

161. Find feasible area for $2x + 3y \leq 450$


➔ To find feasible area for $2x + 3y \leq 450$

Let's draw the line $2x + 3y = 450$ by joining the points $(225, 0)$ & $(0, 150)$




162. Find feasible area for $x - 2y \geq 100$

→ To Find Feasible area for $x - 2y \geq 100$,
Let's draw $x - 2y = 100$ by joining points $(100, 0)$ & $(0, -50)$



163. Find common feasible area for $2x + 3y \leq 60$ & $x, y \geq 0$


→ Let's First draw $2x + 3y = 60$ by joining $(0, 20)$ & $(30, 0)$




$x, y \geq 0$ is known as Non-negativity constraint which restricts feasible area in 1st quadrant.

164. Find common feasible area for $x - 2y \geq 50$ & $2x + 3y \leq 150$

→ Lets draw Lines $x - 2y = 50 : (0, -25) \& (50, 0)$
 $2x + 3y = 150 : (0, 50) \& (75, 0)$



165. Find common feasible area for $x \geq 30, x \leq 60, y \geq 10, y \leq 80$




non-negativity constraint

166. Find common feasible area for $2x + y \leq 60$, $x + y \leq 40$, $x, y \geq 0$

Let's first draw Lines $2x + y = 60$: $(0, 60), (30, 0)$

$x + y = 40$: $(0, 40), (40, 0)$



167. Which of the following point lie in common feasible area of $5x - 3y \leq 80$, $2x + 7y \geq 40$

a. $(5, 2)$

b. $(20, 3)$

~~c. $(30, 30)$~~

d. None

→

$5(5) - 3(2) \leq 80$ ✓
 $2(5) + 7(2) \geq 40$ X

$5(20) - 3(3) \leq 80$ X

$5(30) - 3(30) \leq 80$ ✓
 $2(30) + 7(30) \geq 40$ ✓

↓
 point $(30, 30)$ satisfies both the inequalities

Q: which of the following point lie in common feasible area of $5x + 2y \leq 100$, $8x - 3y \geq 20$, $x + 2y \leq 90$

~~a) $(10, 20)$~~

b) $(80, 60)$

c) Both


d) None

168. Total cost = Fixed cost + Variable cost

Fixed cost : The cost which does not change with change in volume of production is known as Fixed Cost.

Variable cost : The cost which changes in same proportion with change in volume of production is known as Variable Cost.

Semi - Variable cost : If portion of the cost is fixed and portion is variable then cost is said to be semi-variable or semi-fixed cost.



$$y = \text{Total cost}$$

$$x = \text{No. of units of produced}$$

$$F = \text{Fixed cost (Total)}$$

$$V = \text{Vari. cost p.u.}$$

$$y = F + Vx$$

$$\text{Total cost} = \text{Total Fixed cost} + \left(\text{vari. cost p.u.} \times \text{No. of units produced} \right)$$

169. If $b^2 - 4ab = 0$ then roots of quadratic equation are

- a. Equal b. Equal but opposite in sign
 c. Reciprocals of each other ~~d. Can't say~~

Roots are equal when $b^2 - 4ac = 0$
 we can't say anything on the basis of value of $b^2 - 4ab$

170. Sum of 2 numbers is 88 and diff betⁿ first number and half of second number is 10. Find the numbers.

- a. 32,56 b. 44,44 ~~c. 36,52~~ d. 30,58

$$x + y = 88$$

$$x - \frac{y}{2} = 10$$

$$176 - 2y - y = 20$$

$$2x - y = 20$$

$$156 = 3y$$

$$y = 52$$

$$\therefore x = 36$$

$$2(88 - y) - y = 20$$

171. If p, q are roots of $3x^2 - 3x - 1 = 0$. Find value of $(p^3 + q^3)$, $(p^2 + q^2)$

→ ① $p + q = 1$

② $pq = -\frac{1}{3}$

③ $(p^3 + q^3) = (p+q)^3 - 3pq(p+q)$

$= 1^3 - 3(-\frac{1}{3})(1) = 1 + 1 = 2$

④ $(p^2 + q^2) = (p+q)^2 - 2pq = 1^2 - 2(-\frac{1}{3}) = 1 + \frac{2}{3} = \frac{5}{3}$

⑤ $(p-q)^2 = (p+q)^2 - 4pq = 1^2 - 4(-\frac{1}{3}) = 1 + \frac{4}{3} = \frac{7}{3}$

⑥ $p^2q + q^2p = pq(p+q) = -\frac{1}{3} \times 1 = -\frac{1}{3}$

⑦ $\frac{p}{q} + \frac{q}{p} = \frac{p^2 + q^2}{pq} = \frac{5/3}{-1/3} = -5$

⑧ $\frac{p^2}{q} + \frac{q^2}{p} = \frac{p^3 + q^3}{pq} = \frac{2}{-1/3} = -6$

172. If p, q are roots of $3x^2 - 19x - 1 = 0$, whose roots are $\frac{p}{q}$ & $\frac{q}{p}$.

Find quadratic equation.



$$p + q = \frac{19}{3}$$

$$pq = -\frac{1}{3}$$

Quad. eqⁿ whose roots are $\frac{p}{q}$ & $\frac{q}{p}$ is,

$$x^2 - \left(\frac{p}{q} + \frac{q}{p}\right)x + \left(\frac{p}{q} \times \frac{q}{p}\right) = 0$$

$$x^2 - \left(\frac{p^2 + q^2}{pq}\right)x + 1 = 0$$

$$x^2 - \left(\frac{\frac{361}{9} - 2\left(-\frac{1}{3}\right)}{-\frac{1}{3}}\right)x + 1 = 0$$

$$x^2 + \left(\frac{\frac{361}{9} + \frac{6}{9}}{\frac{3}{9}}\right)x + 1 = 0$$

$$x^2 + \frac{367}{3}x + 1 = 0$$

$$3x^2 + 367x + 3 = 0$$

173. The cubic eqⁿ whose roots are m, n, q is :

$$\rightarrow x^3 - (\text{sum of roots})x^2 + [(\text{1st} \times \text{2nd}) + (\text{2nd} \times \text{3rd}) + (\text{1st} \times \text{3rd})]x - (\text{product of roots}) = 0$$

$$x^3 - (m+n+q)x^2 + (mn+nq+mq)x - mnq = 0$$

174. If x = No. of units produced

Fixed Cost = ₹ 3,80,000; Variable Cost p.u. = ₹ 28

then y = Total Cost = _____



Total cost = Total Fixed cost + Total variable cost

$$TC = \text{Total Fixed cost} + (\text{vari. cost p.u.} \times \text{no. of units produced})$$

$$y = 3,80,000 + 28x$$

175. If $(p+2)(p-3) + (p+3)(p-4) = p(2p-5)$, then $p = ?$



$$(p+2)(p-3) + (p+3)(p-4) = p(2p-5)$$

$$p^2 - 3p + 2p - 6 + p^2 - 4p + 3p - 12 = 2p^2 - 5p$$

$$2p^2 - p - 6 - p - 12 = 2p^2 - 5p$$

$$2p^2 - 2p - 18 = 2p^2 - 5p$$

$$2\cancel{p^2} - 2p - 18 - 2\cancel{p^2} + 5p = 0$$

$$3p = 18$$

$$p = 6$$

176. $15x + 23y = -10$ & $3x + 4y = -2$ then $3x + 2y + 2 = ?$



$$\begin{array}{r} 15x + 23y = -10 \\ 15x + 20y = -10 \\ \hline \end{array}$$

$$3y = 0$$

$$y = \frac{0}{3}$$

$$y = 0 \text{ By using calculator.}$$

Let's put $y = 0$ in $15x + 23y = -10$

$$15x + 23(0) = -10$$

$$15x = -10$$

$$x = -\frac{10}{15} = -\frac{2}{3}$$

$$\therefore 3x + 2y + 2 = ?$$

$$= 3\left(-\frac{2}{3}\right) + 2(0) + 2$$

$$= -2 + 0 + 2$$

$$= 0$$



177. Find value of k , if $9x^2 - 24x + k = 0$ has equal roots.

$\rightarrow b^2 - 4ac = 0$
 $(-24)^2 - 4(9)(k) = 0$
 $576 - 36k = 0$
 $576 = 36k$
 $\therefore k = 16$

Roots are equal	$b^2 - 4ac = 0$
Roots are reciprocals of each other	$a = c$
Roots are equal but opposite in sign	$b = 0$

178. Calculate the number such that it is equal to 3 times of its diff from 56.

\rightarrow Let that number be x
 $x = 3 \times (56 - x)$
 $x = 168 - 3x$
 $4x = 168$
 $x = 42$

Roots are	$b^2 - 4ac$
Real, irrational, unequal	$b^2 - 4ac > 0$ & Not a perfect square
Real, Rational, Equal	$b^2 - 4ac = 0$
Real, Rational unequal	$b^2 - 4ac > 0$ & perfect square
Complex/Imagi.	$b^2 - 4ac < 0$

179. $2x + 3y = 5$ & $3x - 4y = 2$ then $5xy = ?$

\rightarrow

$$\begin{array}{r} 6x + 9y = 15 \\ -6x + 8y = -4 \\ \hline 17y = 11 \end{array} \therefore y = \frac{11}{17}$$

$$2x + \left(3 \times \frac{11}{17}\right) = 5$$

$$2x = 5 - \frac{33}{17} = \frac{52}{17} \therefore x = \frac{26}{17}$$

$$5xy = 5 \times \frac{26}{17} \times \frac{11}{17} = \left(\frac{1430}{289}\right)$$

For a cubic eqⁿ
 $ax^3 + bx^2 + cx + d = 0$

Sum of roots
 $= -b/a$

Product of roots
 $= -d/a$

180. $a^2 + b^2 = 45$ then $\frac{1}{a} + \frac{1}{b} = ?$

$ab = 18$

$$\begin{aligned} \rightarrow (a+b)^2 &= (a^2 + b^2) + 2ab & \frac{1}{a} + \frac{1}{b} \\ &= 45 + 2(18) & = \left(\frac{a+b}{ab}\right) \\ (a+b)^2 &= 81 & = \frac{9}{18} = \frac{1}{2} \\ (a+b) &= 9 \end{aligned}$$

181. If roots of quadratic equation are $(2m)$ & $(-2n)$ then factors are :

\rightarrow Roots are : $2m, -2n$
 \therefore Factors are : $(x-2m) \& (x+2n)$

182. If roots of quadratic equation are $(\frac{3}{5})$ & $(\frac{-8}{11})$ then factors are :

\rightarrow Factors are $(5x-3) \& (11x+8)$

Roots of quad. eqn	Quad. eqn
8, 3	$x^2 - 11x + 24 = 0$
11, 9	$x^2 - 20x + 99 = 0$
-6, 88	$x^2 - 82x - 528 = 0$
10, 0	$x^2 - 10x = 0$
$2 + \sqrt{20}, 2 - \sqrt{20}$	$x^2 - 4x - 16 = 0$
$8 + \sqrt{11}, 8 - \sqrt{11}$	$x^2 - 16x + 53 = 0$
0.50, 2.50	$x^2 - 3x + 1.25 = 0, 4x^2 - 12x + 5 = 0$
-6, -9	$x^2 + 15x + 54 = 0$
28, -28	$x^2 - 0x - 784 = 0$



183. If quadratic equation $x^2 - (p+4)x + 2p + 5 = 0$ has equal roots.

Find p

→ AS Roots are equal, $b^2 - 4ac = 0$

$$[-(p+4)]^2 - 4(1)(2p+5) = 0$$

$$p^2 + 8p + 16 - 8p - 20 = 0$$

$$p^2 - 4 = 0$$

$$p^2 = 4$$

$$p = \pm 2$$

184. If $4x^3 + 8x^2 - x - 2 = 0$ then $(2x + 3) = ?$

~~a. 4, -1, 2~~

b. -4, 2, 1

c. 2, -4, -1

d. None

→

$$4x^3 + 8x^2 - x - 2 = 0$$

$$4x^2(x+2) - 1(x+2) = 0$$

$$(x+2)(4x^2 - 1) = 0$$

$$(x+2)[(2x)^2 - 1^2] = 0$$

$$(x+2)(2x-1)(2x+1) = 0$$

$$x = -2, x = \frac{1}{2}, x = -\frac{1}{2}$$

$(2x+3) : 2(-2)+3$	$2 \times \frac{1}{2} + 3$	$2 \times -\frac{1}{2} + 3$
$= -1$	$= 4$	$= 2$

185. Sum of 2 numbers is 15 & their product is 50 then sum of their reciprocal is :

$$\begin{aligned} \rightarrow \quad x + y &= 15 & \frac{1}{x} + \frac{1}{y} \\ xy &= 50 & \\ & & = \frac{x+y}{xy} = \frac{15}{50} = \frac{3}{10} = 0.30 \end{aligned}$$

186. Out of 3 numbers, sum of first and second is 24, sum of 2nd & 3rd is 30, sum of first & third is 26. The smallest number is :

- a. 18 b. 14 c. 16 ~~d. 10~~

\rightarrow 1st Number : x
2nd Number : y
3rd Number : z

$$x + y = 24$$

$$y + z = 30$$

$$x + z = 26$$

$$x + y = 24$$

$$x + 30 - z = 24$$

$$x + 30 - (26 - x) = 24$$

$$x + 30 - 26 + x = 24$$

$$2x + 4 = 24$$

$$2x = 20$$

$$x = 10$$

$$\therefore \boxed{y = 14} \quad \boxed{z = 16}$$

**FORGET THE MISTAKE
REMEMBER THE LESSON !**



A series of horizontal lines for writing notes, consisting of 30 lines in total.

